



# Annals of Representation Theory

SOFIA BRENNER & BURKHARD KÜLSHAMMER 

**Group algebras in which the socle of the center is an ideal**

Volume 1, issue 1 (2024), p. 1-19

<https://doi.org/10.5802/art.1>

Communicated by Radha Kessar.

© The authors, 2024



This article is licensed under the  
CREATIVE COMMONS ATTRIBUTION (CC-BY) 4.0 LICENSE.  
<http://creativecommons.org/licenses/by/4.0/>



NTNU


*Annals of Representation Theory is published by the  
Norwegian University of Science and Technology  
and a member of the  
Centre Mersenne for Open Scientific Publishing*

e-ISSN: pending





# Group algebras in which the socle of the center is an ideal

Sofia Brenner and Burkhard Külshammer\* 

**ABSTRACT.** Let  $F$  be a field of characteristic  $p > 0$ . We study the structure of the finite groups  $G$  for which the socle of the center of  $FG$  is an ideal in  $FG$  and classify the finite  $p$ -groups  $G$  with this property. Moreover, we give an explicit description of the finite groups  $G$  for which the Reynolds ideal of  $FG$  is an ideal in  $FG$ .

## 1. INTRODUCTION

Let  $F$  be a field and consider the group algebra  $FG$  of a finite group  $G$  and its center  $ZFG$ . The question when the Jacobson radical of  $ZFG$  is an ideal in  $FG$  has been answered by Clarke [4], Koshitani [7] and Külshammer [9]. We now study the corresponding problem for the socle  $\text{soc}(ZFG)$  of  $ZFG$  as well as for the Reynolds ideal  $R(FG)$  of  $FG$ . In a prequel to this paper [3], we have already given some approaches to these problems for general symmetric algebras. Now, our aim is to analyze the structure of the finite groups  $G$  for which  $\text{soc}(ZFG)$  or  $R(FG)$  are ideals of  $FG$  in a group-theoretic manner. For the Reynolds ideal, we obtain the following characterization:

**Theorem A.** *Let  $F$  be a field of characteristic  $p > 0$  and let  $G$  be a finite group. Then the Reynolds ideal  $R(FG)$  is an ideal in  $FG$  if and only if  $G'$  is contained in the  $p$ -core  $O_p(G)$  of  $G$ .*

As a consequence of this result, it follows that if  $\text{soc}(ZFG)$  is an ideal in  $FG$ , one has  $G = P \rtimes H$  for a Sylow  $p$ -subgroup  $P$  of  $G$  and an abelian  $p'$ -group  $H$ . Based on this decomposition, we derive some fundamental results on the structure of finite groups  $G$  for which  $\text{soc}(ZFG)$  is an ideal in  $FG$ . Subsequently, we classify the finite  $p$ -groups  $G$  with this property:

**Theorem B.** *Let  $F$  be a field of characteristic  $p > 0$  and let  $G$  be a finite  $p$ -group. Then  $\text{soc}(ZFG)$  is an ideal in  $FG$  if and only if*

---

Manuscript received 2022-12-02 and accepted 2023-06-07.

2020 *Mathematics Subject Classification.* 20C05, 20C20, 16S34.

\* Corresponding author.

- (i)  $G$  has nilpotency class at most two, that is,  $G' \subseteq Z(G)$  holds, or  
(ii)  $p = 2$  and  $G' \subseteq Y(G)Z(G)$  with  $Y(G) = \langle fg^{-1} : \{f, g\} \text{ is a conjugacy class of length } 2 \text{ of } G \rangle$ .

In particular,  $G$  is metabelian.

Note that since the  $p$ -groups of nilpotency class at most two form a large subclass of the finite  $p$ -groups, the condition that  $\text{soc}(ZFG)$  is an ideal in  $FG$  is often satisfied. One implication of Theorem B generalizes to arbitrary finite groups:

**Theorem C.** *Let  $F$  be a field of characteristic  $p > 0$  and let  $G$  be a finite group. Suppose that one of the following holds:*

- (i)  $G' \subseteq Z(O_p(G))$ , or  
(ii)  $p = 2$  and  $G' \subseteq Y(O_p(G))Z(O_p(G))$ .

Then  $\text{soc}(ZFG)$  is an ideal in  $FG$ .

The above results are major ingredients for the proof of the main result of this paper, which is a decomposition of  $G$  into a central product:

**Theorem D.** *Let  $F$  be a field of characteristic  $p > 0$ . Suppose that  $G$  is a finite group for which  $\text{soc}(ZFG)$  is an ideal in  $FG$  and write  $G = P \rtimes H$  for a Sylow  $p$ -subgroup  $P$  of  $G$  and an abelian  $p'$ -group  $H$  as before. Then  $G$  is the central product of the centralizer  $C_P(H)$  and the  $p$ -residual group  $O^p(G)$ . Moreover,  $\text{soc}(ZFC_P(H))$  and  $\text{soc}(ZFO^p(G))$  are ideals in  $FC_P(H)$  and  $FO^p(G)$ , respectively. Furthermore, we have*

$$\text{soc}(ZFG) = (Z(P)G')^+ \cdot FG,$$

where  $(Z(P)G')^+ \in FG$  denotes the sum of the elements in  $Z(P)G'$ .

This statement will allow us to restrict our investigation to the case  $P = G'$ . A detailed analysis of the structure of finite groups  $G$  for which  $\text{soc}(ZFG)$  is an ideal in  $FG$ , based on the above results, will be carried out in a sequel to this paper.

We proceed as follows: First, we introduce our notation (see Section 2) and study the general structure of the finite groups  $G$  for which  $\text{soc}(ZFG)$  or  $R(FG)$  are ideals in  $FG$  (see Section 3). In Section 4, we classify the  $p$ -groups  $G$  for which  $\text{soc}(ZFG)$  is an ideal in  $FG$  for a field  $F$  of characteristic  $p > 0$ . In Section 5, we derive the decomposition of  $G$  given in Theorem D.

## 2. NOTATION

Let  $G$  be a finite group and  $p$  a prime number. As customary, let  $G'$ ,  $Z(G)$  and  $\Phi(G)$  denote the derived subgroup, the center and the Frattini subgroup of  $G$ , respectively. For elements  $a, b \in G$ , we define their commutator as  $[a, b] = aba^{-1}b^{-1}$ . We write  $[g]$  for the conjugacy class of  $g \in G$  and set  $\text{Cl}(G)$  to be the set of conjugacy classes of  $G$ . The nilpotency class of a nilpotent group  $G$  will be denoted by  $c(G)$ . Recall that every  $p$ -group is nilpotent. For subsets  $S$  and  $T$  of  $G$ , let  $C_T(S)$  and  $N_T(S)$  denote the centralizer and the normalizer of  $S$  in  $T$ , respectively. As customary, let  $O_p(G)$ ,  $O_{p'}(G)$  and  $O_{p',p}(G)$  be the  $p$ -core, the  $p'$ -core and the  $p', p$ -core of  $G$ , respectively. By  $O^p(G)$  and  $O^{p'}(G)$ , we denote the  $p$ -residual subgroup and the  $p'$ -residual subgroup of  $G$ , respectively. As customary, let  $g_p$  and  $g_{p'}$  be the  $p$ -part and the  $p'$ -part of an element  $g \in G$ , respectively. The  $p'$ -section of  $g$  is given by all elements in  $G$  whose  $p'$ -part is conjugate to  $g_{p'}$ . We write  $G = G_1 * G_2$  if  $G$  is the central product of subgroups  $G_1$  and  $G_2$ , that is, we have  $G = \langle G_1, G_2 \rangle$  and  $[G_1, G_2] = 1$ .

For a field  $F$  and a finite-dimensional  $F$ -algebra  $A$ , we denote by  $J(A)$  and  $\text{soc}(A)$  its Jacobson radical and (left) socle, the sum of all minimal left ideals of  $A$ , respectively. Both  $J(A)$  and  $\text{soc}(A)$  are ideals in  $A$ . In this paper, an ideal  $I$  of  $A$  is always meant to be a two-sided ideal, and we denote it by  $I \trianglelefteq A$ . Additionally, we study the Reynolds ideal  $R(A) := \text{soc}(A) \cap Z(A)$  of  $A$ . Furthermore, let  $K(A)$  denote the commutator space of  $A$ , that is, the  $F$ -subspace of  $A$  spanned by all elements of the form  $ab - ba$  with  $a, b \in A$ .

In the following, we consider the group algebra  $FG$  of  $G$  over  $F$ . Recall that  $FG$  is a symmetric algebra with symmetrizing linear form

$$\lambda: FG \rightarrow F, \quad \sum_{g \in G} a_g g \mapsto a_1. \quad (2.1)$$

For subsets  $S$  and  $T$  of  $FG$ , we write  $\text{lAnn}_T(S)$  and  $\text{rAnn}_T(S)$  for the left and the right annihilator of  $S$  in  $T$ , respectively, and  $\text{Ann}_T(S)$  if both subspaces coincide. For  $H \subseteq G$ , we set  $H^+ := \sum_{h \in H} h \in FG$ . It is well-known that the elements  $C^+$  with  $C \in \text{Cl}(G)$  form an  $F$ -basis of the center  $ZFG$  of  $FG$ .

In this paper, we mainly study the Jacobson radical  $J(ZFG)$  and the socle  $\text{soc}(ZFG)$  of the center of  $FG$  as well as the Reynolds ideal  $R(FG)$ . All three spaces are ideals in  $ZFG$ , but not necessarily in  $FG$ . Note that  $J(ZFG) = J(FG) \cap ZFG$  holds (see [10, Theorem 1.10.8]) and that by [10, Theorem 1.10.22], we have  $\text{soc}(ZFG) = \text{Ann}_{ZFG}(J(ZFG))$ . Furthermore, observe that  $J(ZFG)$ ,  $\text{soc}(ZFG)$  and  $R(FG)$  are ideals in  $FG$  if and only if they are closed under multiplication with elements of  $FG$  since they are additively closed.

We recall the definition of the augmentation ideal

$$\omega(FG) = \left\{ \sum_{g \in G} a_g g \in FG : \sum_{g \in G} a_g = 0 \right\}.$$

An  $F$ -basis of  $\omega(FG)$  is given by  $\{1 - g : 1 \neq g \in G\}$ . If  $F$  is a field of characteristic  $p > 0$  and  $G$  is a  $p$ -group, then  $J(FG)$  and  $\omega(FG)$  coincide (see [10, Theorem 1.11.1]). For a normal subgroup  $N$  of  $G$ , we consider the canonical projection

$$\nu_N: FG \rightarrow F[G/N], \quad \sum_{g \in G} a_g g \mapsto \sum_{g \in G} a_g \cdot gN.$$

Its kernel is given by  $\omega(FN) \cdot FG = FG \cdot \omega(FN)$  (see [10, Proposition 1.6.4]).

### 3. GENERAL PROPERTIES

Let  $F$  be a field. In this part, we answer the question for which finite groups  $G$  the Reynolds ideal  $R(FG)$  is an ideal in  $FG$ . Moreover, we derive structural results on finite groups  $G$  for which  $\text{soc}(ZFG)$  is an ideal in  $FG$ . In the next section, these will be applied in order to classify the finite groups of prime power order with this property.

Concerning the choice of the underlying field  $F$ , we note the following:

#### Remark 3.1.

- (i) Assume that  $F$  is of characteristic zero or of positive characteristic not dividing  $|G|$ . By Maschke's theorem, the group algebra  $FG$  is semisimple. In particular,  $J(FG) = J(ZFG) = 0$  follows, which yields  $R(FG) = \text{soc}(ZFG) = ZFG$ . Since  $FG$  is unitary,  $\text{soc}(ZFG)$  is an ideal of  $FG$  if and only if  $ZFG = FG$  holds, that is, if and only if  $G$  is abelian.

- (ii) Let  $F$  be a field of characteristic  $p > 0$  and let  $G$  be a finite group. Then  $\text{soc}(Z\mathbb{F}_p G)$  is an ideal in  $\mathbb{F}_p G$  if and only if  $\text{soc}(ZFG)$  is an ideal in  $FG$ . A similar statement holds for the Reynolds ideal.

From now on until the end of this paper, we therefore assume that  $F$  is an algebraically closed field of characteristic  $p > 0$ .

This section is organized as follows: We first derive a criterion for  $\text{soc}(ZFG) \trianglelefteq FG$  (see Section 3.1) and answer the question when the Reynolds ideal of  $FG$  is an ideal in  $FG$  (see Section 3.2). In Section 3.3, we investigate  $p$ -blocks of  $FG$ . Subsequently, we find a basis for  $J(ZFG)$  (see Section 3.4) and construct elements in  $\text{soc}(ZFG)$  arising from normal  $p$ -subgroups of  $G$  (see Section 3.5). In Section 3.6, we study the case that  $G'$  is contained in the center of a Sylow  $p$ -subgroup of  $G$ . We conclude this part by investigating the transition to quotient groups in Section 3.7 and studying central products in Section 3.8.

**3.1. Criterion for  $\text{soc}(ZFG) \trianglelefteq FG$ .** Let  $G$  be a finite group. In this section, we derive an equivalent criterion for  $\text{soc}(ZFG) \trianglelefteq FG$ .

**Lemma 3.2.** *We have  $FG \cdot K(FG) = FG \cdot \omega(FG')$ .*

*Proof.* As  $FG/\omega(FG') \cdot FG$  is isomorphic to the commutative algebra  $F[G/G']$ , we have  $K(FG) \subseteq \omega(FG') \cdot FG$  and hence  $K(FG) \cdot FG \subseteq \omega(FG') \cdot FG$  follows. Now let  $f: FG \rightarrow FG/K(FG) \cdot FG$  be the canonical projection map. For all  $a, b \in G$ , we have  $f([a, b]) = f(a)f(b)f(a)^{-1}f(b)^{-1} = 1$  since  $FG/K(FG) \cdot FG$  is a commutative algebra. For  $g \in G'$ , this yields  $f(g) = 1$  and hence  $f(g - 1) = 0$ . This shows  $\omega(FG') \subseteq \text{Ker}(f) = K(FG) \cdot FG$ , which proves the claim.  $\square$

**Lemma 3.3.** *The socle  $\text{soc}(ZFG)$  is an ideal in  $FG$  if and only if  $\text{soc}(ZFG) \subseteq (G')^+ \cdot FG$  holds.*

*Proof.* By [9, Lemma 2.1], we have  $\text{soc}(ZFG) \trianglelefteq FG$  if and only if  $K(FG) \cdot \text{soc}(ZFG) = 0$  holds, which is equivalent to  $FG \cdot K(FG) \cdot \text{soc}(ZFG) = 0$ . By Lemma 3.2, this is equivalent to  $FG \cdot \omega(FG') \cdot \text{soc}(ZFG) = 0$ , that is, to  $\text{soc}(ZFG) \subseteq \text{rAnn}_{FG}(\omega(FG')) = (G')^+ \cdot FG$  (see [11, Lemma 3.1.2]).  $\square$

**3.2. Reynolds ideal.** Let  $G$  be a finite group. In this section, we answer the question when the Reynolds ideal  $R(FG)$  is an ideal in  $FG$ . Our main result is the following:

**Theorem 3.4.** *The following properties are equivalent:*

- (i)  $R(FG)$  is an ideal of  $FG$ .
- (ii)  $G' \subseteq O_p(G)$ .
- (iii)  $G = P \rtimes H$  with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$ .

*In this case, we have  $R(FG) = O_p(G)^+ \cdot FG$ .*

*Proof.* Suppose that  $R(FG)$  is an ideal in  $FG$ . Then  $FG$  is a basic  $F$ -algebra by [3, Lemma 2.2]. Since  $F$  is algebraically closed, this implies that  $FG/J(FG)$  is commutative. By Lemma 3.2, we have  $\omega(FG') \cdot FG = K(FG) \cdot FG \subseteq J(FG)$ . Thus, for  $g \in G'$ , the element  $g - 1$  is nilpotent. Hence there exists  $n \in \mathbb{N}$  with  $0 = (g - 1)^{p^n} = g^{p^n} - 1$ . This shows that  $G'$  is a  $p$ -group and hence contained in  $O_p(G)$ .

Now assume  $G' \subseteq O_p(G)$  and let  $P \in \text{Syl}_p(G)$ . Then  $G' \subseteq P$  follows, so  $P$  is a normal subgroup of  $G$  and  $G/P$  is abelian. By the Schur-Zassenhaus theorem,  $P$  has a complement  $H$  in  $G$ . Moreover,  $H$  is isomorphic to  $G/P$  and thus abelian.

Finally suppose that  $G = P \rtimes H$  holds, where  $P \in \text{Syl}_p(G)$  and  $H$  is an abelian  $p'$ -group. In particular, we have  $P = O_p(G)$ . We obtain  $J(FG) = \omega(FP) \cdot FG$  and  $\text{soc}(FG) = \text{Ann}_{FG}(J(FG)) = P^+ \cdot FG \subseteq (G')^+ \cdot FG \subseteq ZFG$ , so that  $R(FG) = P^+ \cdot FG$  is an ideal in  $FG$ .  $\square$

This proves Theorem A. Moreover, we obtain the following necessary condition for  $\text{soc}(ZFG) \trianglelefteq FG$ :

**Corollary 3.5.** *If  $\text{soc}(ZFG)$  is an ideal of  $FG$ , we have  $G = P \rtimes H$  with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$ .*

*Proof.* By [3, Lemma 1.3],  $\text{soc}(ZFG) \trianglelefteq FG$  implies  $R(FG) \trianglelefteq FG$ . With this, the claim follows from Theorem 3.4.  $\square$

**Remark 3.6.** Let  $G = P \rtimes H$  with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$ .

- (i) By [5, Theorem 5.3.5], we have  $P = C_P(H)[P, H]$ . Due to  $[P, H] \subseteq G'$ , this yields  $G = HP = HC_P(H)[P, H] = HC_P(H)G'$ . Note that  $[G, H] = [P, H] = [[P, H], H] = [G', H] = [[G', H], H]$  holds by [5, Theorem 5.3.6] and that this is a normal subgroup of  $PH = G$ .
- (ii) We have  $O^p(G) = N$  for  $N := H[G, H]$ : Clearly,  $N$  is a normal subgroup of  $G$ . Since  $G/N$  is a  $p$ -group, we have  $O^p(G) \subseteq N$ . On the other hand,  $G/O^p(G)$  is a  $p$ -group, which implies  $H \subseteq O^p(G)$  and hence  $N \subseteq O^p(G)$  as  $O^p(G)$  is a normal subgroup of  $G$ . In particular, this implies  $O^p(G)' \subseteq [G, H]$ . On the other hand, we have  $[G, H] = [[G', H], H] \subseteq [O^p(G), O^p(G)] = O^p(G)'$  by (i) and hence  $O^p(G)' = [G, H] \in \text{Syl}_p(O^p(G))$  follows.
- (iii) Since  $O_{p'}(G)$  is contained in the abelian group  $H$  and  $[P, O_{p'}(G)] \subseteq P \cap O_{p'}(G) = 1$  holds, we have  $O_{p'}(G) \subseteq Z(G)$ . Hence [5, Theorem 6.3.3] implies  $C_G(P) \subseteq O_{p'}(G) = O_{p'}(G) \times P$ , and we conclude that  $C_G(P) = O_{p'}(G) \times Z(P)$  holds.
- (iv) Since  $R(FG)$  is spanned by the  $p'$ -section sums of  $G$  (see [8, Equation (39)]), every  $p'$ -section is of the form  $hP$  for some  $h \in H$ .

**3.3. Blocks and the  $p'$ -core.** Let  $G$  be an arbitrary finite group. In this section, we investigate the conditions  $\text{soc}(Z(B)) \trianglelefteq B$  and  $R(B) \trianglelefteq B$  for a  $p$ -block  $B$  of  $FG$ .

**Remark 3.7.** Let  $FG = B_1 \oplus \dots \oplus B_n$  be the decomposition of  $FG$  into its  $p$ -blocks. Then we have

$$\text{soc}(ZFG) = \text{soc}(Z(B_1)) \oplus \dots \oplus \text{soc}(Z(B_n)).$$

In particular,  $\text{soc}(ZFG)$  is an ideal in  $FG$  if and only if  $\text{soc}(Z(B_i)) \trianglelefteq B_i$  holds for all  $i \in \{1, \dots, n\}$ , and the analogous statement is true for the Reynolds ideal. Furthermore, it is known that the principal blocks of  $FG$  and  $F\bar{G}$  are isomorphic for  $\bar{G} := G/O_{p'}(G)$ .

For the Reynolds ideal, we obtain the following result:

**Lemma 3.8.** *The following are equivalent:*

- (i) *There exists a block  $B$  of  $FG$  for which  $R(B) \trianglelefteq B$  holds.*
- (ii) *For the principal block  $B_0$  of  $FG$ , we have  $R(B_0) \trianglelefteq B_0$ .*
- (iii)  *$G'$  is contained in  $O_{p'}(G)$ .*

*Proof.* Assume that (i) holds. By [9, Proposition 4.1], this implies  $B \cong B_0$  and hence (ii) holds. Now assume that (ii) holds. By [9, Remarks 2.2 and 3.1], every simple  $B_0$ -module is one-dimensional. Since the intersection of the kernels of the simple  $B_0$ -modules is given by  $O_{p'}(G)$  (see [1, Theorem 2]), we obtain  $G' \subseteq O_{p'}(G)$ . Finally, assume that (iii) holds.

Then we have  $\bar{G}' \subseteq O_p(\bar{G})$ . Theorem 3.4 yields  $R(F\bar{G}) \trianglelefteq F\bar{G}$ , which implies  $R(\bar{B}_0) \trianglelefteq \bar{B}_0$  by Remark 3.7. Since  $B_0$  and  $\bar{B}_0$  are isomorphic, we obtain  $R(B_0) \trianglelefteq B_0$ .  $\square$

Concerning the analogous problem for the socle of the center, we first observe the following:

**Lemma 3.9.** *The following are equivalent:*

- (i) *There exists a block  $B$  of  $FG$  for which  $\text{soc}(Z(B)) \trianglelefteq B$  holds.*
- (ii) *For the principal block  $B_0$  of  $FG$ , we have  $\text{soc}(Z(B_0)) \trianglelefteq B_0$ .*
- (iii) *For the principal block  $\bar{B}_0$  of  $F\bar{G}$ , we have  $\text{soc}(Z(\bar{B}_0)) \trianglelefteq \bar{B}_0$ .*

*Proof.* As in the proof of Lemma 3.8, the equivalence of (i) and (ii) follows by [9, Proposition 4.1] and the equivalence of (ii) and (iii) follows from the fact that  $B_0$  and  $\bar{B}_0$  are isomorphic.  $\square$

This has the following important consequence:

**Lemma 3.10.** *We have  $\text{soc}(ZFG) \trianglelefteq FG$  if and only if  $R(FG) \trianglelefteq FG$  and  $\text{soc}(ZF\bar{G}) \trianglelefteq F\bar{G}$  hold.*

*Proof.* If  $\text{soc}(ZFG)$  is an ideal of  $FG$ , then  $R(FG) \trianglelefteq FG$  holds by [3, Lemma 1.3] and  $\text{soc}(ZF\bar{G})$  is an ideal of  $F\bar{G}$  by [3, Proposition 2.10]. For the latter, note that  $F\bar{G} \cong FG/\text{Ker}(\nu_{O_{p'}(G)})$  can be viewed as a quotient algebra of  $FG$ . Now let  $R(FG)$  and  $\text{soc}(ZF\bar{G})$  be ideals in  $FG$  and  $F\bar{G}$ , respectively. By Remark 3.7, this yields  $\text{soc}(Z(\bar{B}_0)) \trianglelefteq \bar{B}_0$  and hence  $\text{soc}(Z(B_0)) \trianglelefteq B_0$  (see Lemma 3.9). Since  $R(FG)$  is an ideal in  $FG$ , all blocks of  $FG$  are isomorphic to  $B_0$  by [9, Proposition 4.1]. By Remark 3.7, we then obtain  $\text{soc}(ZFG) \trianglelefteq FG$ .  $\square$

**Remark 3.11.** Assume that  $G$  is of the form  $G = P \rtimes H$  with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$ . Then  $\text{soc}(ZFG) \trianglelefteq FG$  is equivalent to  $\text{soc}(ZF\bar{G}) \trianglelefteq F\bar{G}$  (see Theorem 3.4 and Lemma 3.10). By going over to the quotient group  $G/O_{p'}(G)$ , we may therefore restrict our investigation to groups  $G$  with  $O_{p'}(G) = 1$ .

**3.4. Basis for  $J(ZFG)$ .** Let  $G = P \rtimes H$  be a finite group with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$  (see Theorem 3.4). The aim of this section is to determine an  $F$ -basis for  $J(ZFG)$ . In the given situation, the kernel of the canonical map  $\nu_P: FG \rightarrow F[G/P]$  is given by  $J(FG)$  (see [10, Corollary 1.11.11]). In the following, we distinguish two types of conjugacy classes:

**Remark 3.12.** Let  $C \in \text{Cl}(G)$ . We obtain  $|\bar{C}| = 1$  for the image  $\bar{C} \in \text{Cl}(G/P)$  of  $C$  in  $G/P$  since this group is abelian. Now two cases can occur:

- $|C|$  is divisible by  $p$ : Then  $\nu_P(C^+) = |C| \cdot \bar{C}^+ = 0$  yields  $C^+ \in \text{Ker}(\nu_P) \cap ZFG = J(ZFG)$ .
- $|C|$  is not divisible by  $p$ : In this case,  $|P|$  divides  $|C_G(g)|$  for any  $g \in C$ . This yields  $P \subseteq C_G(g)$  and hence  $C \subseteq C_G(P)$ . As customary, we decompose  $g = g_{p'}g_p$  into its  $p'$ -part and  $p$ -part. Note that  $g_{p'} \in O_{p'}(G) \subseteq Z(G)$  and  $g_p \in Z(P)$  hold by Remark 3.6. Due to  $g_{p'} \in Z(G)$ , we have  $C = g_{p'}[g_p]$  and the element  $C^+ - |C| \cdot g_{p'}$  is contained in  $\text{Ker}(\nu_P) \cap ZFG = J(ZFG)$ .

**Definition 3.13.** For  $C \in \text{Cl}(G)$  with  $C \not\subseteq O_{p'}(G)$ , we set  $b_C := C^+$  if  $p$  divides  $|C|$ , and  $b_C := C^+ - |C| \cdot g_{p'}$  otherwise.

With this, we obtain the following basis for  $J(ZFG)$ :



**Theorem 3.14.** *An  $F$ -basis for  $J(ZFG)$  is given by  $B := \{b_C : C \in \text{Cl}(G), C \not\subseteq O_{p'}(G)\}$ .*

*Proof.* By Remark 3.12, we have  $B \subseteq J(ZFG)$ . Note that the elements in  $B \cup O_{p'}(G)$  form an  $F$ -basis for  $ZFG$ . Since the algebra  $FO_{p'}(G)$  is semisimple,  $J(ZFG)$  is spanned by  $B$ . □

**Remark 3.15.** The decomposition  $FG = \bigoplus_{h \in H} FhP$  gives rise to an  $H$ -grading of  $FG$ . Note that the basis of  $J(ZFG)$  given in Theorem 3.14 consists of homogeneous elements with respect to this grading. In particular,  $J(ZFG)$  is a  $H$ -graded subspace of  $FG$ . It follows that  $\text{soc}(ZFG) = \text{Ann}_{ZFG}(J(ZFG))$  is a  $H$ -graded subspace of  $FG$  as well, that is, we have

$$\text{soc}(ZFG) = \bigoplus_{h \in H} (\text{soc}(ZFG) \cap FhP).$$

**3.5. Elements in  $\text{soc}(ZFG)$ .** Let  $G$  be an arbitrary finite group. In this section, we study elements of  $\text{soc}(ZFG)$  which arise from certain normal  $p$ -subgroups of  $G$ . Using these, we show that  $G'$  has nilpotency class at most two if  $\text{soc}(ZFG)$  is an ideal in  $FG$ . Moreover, we derive a decomposition of  $G$  which will later be used to prove Theorem D.

**Lemma 3.16.** *Let  $N$  be a normal  $p$ -subgroup of  $G$  and set  $M := \{x \in [N, G] : x^p \in [N, [N, G]]\}$ . For  $C \in \text{Cl}(G)$  with  $C \not\subseteq C_G(N)$ , we have  $\nu_M(C^+) = 0$  and hence  $M^+ \cdot C^+ = 0$ . In particular, this implies  $\nu_N(C^+) = 0$  and  $N^+ \cdot C^+ = 0$ .*

*Proof.* Note that  $M$  is a normal subgroup of  $G$ . Let  $R$  be an orbit of the conjugation action of  $N$  on  $C$  and consider an element  $r \in R$ . Then  $C \not\subseteq C_G(N)$  implies  $N \not\subseteq C_G(r)$ , which yields  $|R| = |N : C_N(r)| \neq 1$ . Set  $X := \langle N, R \rangle = \langle N, r \rangle$ .

First consider the case  $[N, G] \subseteq Z(N)$ . Then the map  $f : N \rightarrow N, n \mapsto [n, r]$  is a group endomorphism with kernel  $C_N(r)$ . We set  $S := \text{Im}(f)$ . Then we have  $|R| = |N : C_N(r)| = |S|$ , so in particular,  $|S|$  is a nontrivial power of  $p$ . Let  $\tilde{G} := G/M$  and set  $\tilde{g} := gM \in \tilde{G}$  for  $g \in G$  (similarly for subsets of  $G$ ). Note that  $\tilde{R}$  is an orbit of the conjugation action of  $\tilde{N}$  on  $\tilde{C}$ . As before, we obtain  $|\tilde{R}| = |\tilde{N} : C_{\tilde{N}}(\tilde{r})| = |\tilde{S}| = |S : S \cap M|$ . Since  $S \subseteq [N, G]$  is a nontrivial  $p$ -group,  $|S \cap M|$  is divisible by  $p$ . With this, we obtain

$$\nu_M(R^+) = \frac{|R|}{|\tilde{R}|} \cdot \tilde{R}^+ = |S \cap M| \cdot \tilde{R}^+ = 0.$$

Now we consider the general case. Let  $L := [N, [N, G]]$ . We set  $\tilde{G} := G/L$  and write  $\tilde{g} := gL \in \tilde{G}$  for  $g \in G$  (similarly for subsets of  $G$ ). Note that we have  $[\tilde{N}, [\tilde{N}, \tilde{G}]] = 1$  and hence  $[\tilde{N}, \tilde{G}] \subseteq Z(\tilde{N})$ . First assume  $C_{\tilde{N}}(\tilde{r}) = \tilde{N}$ . For any  $n \in N$ , one then has  $[n, r] \in L$ , which implies  $\nu_L(R^+) = |R| \cdot \tilde{r} = 0$ . Due to  $L \subseteq M$ , this yields  $\nu_M(R^+) = 0$ . Now assume  $C_{\tilde{N}}(\tilde{r}) \subsetneq \tilde{N}$ . In particular, we have  $\tilde{C} \not\subseteq C_{\tilde{G}}(\tilde{N})$ . The first part of the proof yields  $\nu_{\tilde{M}}(\tilde{R}^+) = 0$ , which implies

$$\nu_{\tilde{M}}(\nu_L(R^+)) = \nu_{\tilde{M}}\left(\frac{|R|}{|\tilde{R}|} \cdot \tilde{R}^+\right) = \frac{|R|}{|\tilde{R}|} \cdot \nu_{\tilde{M}}(\tilde{R}^+) = 0.$$

Due to  $\tilde{G}/\tilde{M} = (G/L)/(M/L) \cong G/M$ , the map  $\nu_{\tilde{M}} \circ \nu_L$  can be identified with  $\nu_M$  and hence  $\nu_M(R^+) = 0$  follows. Since  $R$  was arbitrary, this yields  $\nu_M(C^+) = 0$ . In particular, we have  $M^+ \cdot C^+ = 0$ . □

**Proposition 3.17.** *Let  $N$  be a normal  $p$ -subgroup of  $G$  and set  $M := \{x \in [N, G] : x^p \in [N, [N, G]]\}$  as in Lemma 3.16. Moreover, let  $K$  be a characteristic subgroup of  $C_G(N)$  which satisfies  $K^+ \in \text{soc}(ZFC_G(N))$ . Then we have  $(MK)^+ \in \text{soc}(ZFG)$ . In particular, this applies to  $K := O^{p'}(C_G(N))$ .*

*Proof.* By Lemma 3.16,  $ZFG$  is the sum of the subspaces  $ZFG \cap FC_G(N)$  and  $ZFG \cap \text{Ker}(\nu_M)$ . Since  $\text{Ker}(\nu_M) = \omega(FM)FG = J(FM)FG \subseteq J(FG)$  holds (see [10, Proposition 1.6.4]), we have  $ZFG \cap \text{Ker}(\nu_M) \subseteq J(ZFG)$ . Since  $ZFG \cap FC_G(N)$  is contained in  $ZFC_G(N)$ , the space  $J(ZFG \cap FC_G(N)) \subseteq J(ZFC_G(N))$  is annihilated by  $K^+$ . This proves that  $(MK)^+$  annihilates  $J(ZFG)$ . Now let  $K := O^{p'}(C_G(N))$ . Since  $K^+$  annihilates  $J(FC_G(N)) = J(FK)FC_G(N)$  (see [10, Theorem 1.11.10]), we have  $K^+ \in \text{soc}(ZFC_G(N))$  as required.  $\square$

Now we return to the assumption that  $G$  is of the form  $P \rtimes H$  with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$  as in Theorem 3.4.

**Lemma 3.18.** *Suppose that  $N$  is a normal  $p$ -subgroup of  $G$ . Then  $(C_P(N)M)^+ \in \text{soc}(ZFG)$  follows, where  $M := \{x \in [N, G] : x^p \in [N, [N, G]]\}$  is defined as in Lemma 3.16. In particular, we have  $(C_P(N)N)^+ \in \text{soc}(ZFG)$ . If  $\text{soc}(ZFG)$  is an ideal in  $FG$ , then  $G' \subseteq C_P(N)M$  follows.*

*Proof.* Since  $C_P(N)$  is a normal Sylow  $p$ -subgroup of  $C_G(N)$ , we have  $O^{p'}(C_G(N)) = C_P(N)$ . Proposition 3.17 then yields  $(C_P(N)M)^+ \in \text{soc}(ZFG)$ . Since  $C_P(N)N$  is a union of cosets of  $C_P(N)M$ , we obtain  $(C_P(N)N)^+ \in \text{soc}(ZFG)$ . If  $\text{soc}(ZFG)$  is an ideal in  $FG$ , then  $G' \subseteq C_P(N)M$  follows by Lemma 3.3.  $\square$

The following result will be particularly useful for our derivation on  $p$ -groups:

**Corollary 3.19.** *We have  $(Z(P)G')^+ \cdot FG \subseteq \text{soc}(ZFG) \subseteq O_p(Z(G))^+ \cdot FG$ .*

*Proof.* By Lemma 3.18, we obtain  $(Z(P)M)^+ \in \text{soc}(ZFG)$  for  $M = \{x \in [P, G] : x^p \in [P, [P, G]]\} \subseteq G'$ . In particular, this implies  $(Z(P)G')^+ \in \text{soc}(ZFG)$ . Since we have  $(Z(P)G')^+ \cdot FG \subseteq (G')^+ \cdot FG \subseteq ZFG$ , this implies  $(Z(P)G')^+ \cdot FG \subseteq \text{soc}(ZFG)$ . Now for  $z \in O_p(Z(G))$ , the element  $z - 1$  is nilpotent and hence contained in  $J(ZFG)$ . For  $x = \sum_{g \in G} a_g g \in \text{soc}(ZFG)$ , this yields  $x \cdot (z - 1) = 0$ , which translates to  $a_g = a_{gz}$  for all  $g \in G$ . Hence  $x \in O_p(Z(G))^+ \cdot FG$  follows.  $\square$

Observe that the right inclusion in the preceding lemma holds for arbitrary finite groups. The next result is the central ingredient in the proof of Theorem D:

**Proposition 3.20.** *Suppose that  $G' \subseteq C_P(N)N$  holds for every normal  $p$ -subgroup  $N$  of  $G$ . Then the following hold:*

- (i) *We have  $[P, G'] \subseteq Z(G')$ . In particular, this implies  $G'' \subseteq Z(P)$  and that the nilpotency class of  $G'$  is at most two. Moreover, we obtain  $\Phi(G') \subseteq Z(G')$ .*
- (ii) *We have  $P = C_P(H) * [P, H]$  and  $G = C_P(H) * O^p(G)$ .*

*Proof.*

- (i) Let  $D$  be a critical subgroup of  $P$  (in the sense of [5, Theorem 5.3.11]). Then  $D$  is normal in  $G$ , and  $Z(D)$  contains  $\Phi(D)$ ,  $C_P(D)$  and  $[P, D]$ . By assumption, we have  $G' \subseteq DC_P(D) = D$ . Hence we have

$$[P, G'] \subseteq [P, D] \subseteq Z(D) \subseteq C_G(G'),$$

which implies  $[P, G'] \subseteq Z(G')$ . With the 3-subgroups lemma, we obtain  $[G'', P] = [[G', G'], P] = 1$ , that is,  $G'' \subseteq Z(P)$ . Furthermore, for  $x \in G'$ , we have  $x \in D$  and hence  $x^p \in Z(D) \subseteq C_G(G')$ , which implies  $x^p \in Z(G')$ .

- (ii) By (i), we have  $B := [C_P(H), [P, H]] \subseteq [P, G'] \subseteq Z(G')$ . Furthermore,  $B$  is normal in  $C_P(H)[P, H] = P$  and  $PH = G$ . Due to

$$[C_P(H), G] = [C_P(H), C_P(H)[P, H]H] = [C_P(H), C_P(H)[P, H]] \subseteq C_P(H)B,$$

the subgroup  $N := C_P(H)B$  is normal in  $G$ . Moreover, we find  $[N, H] = [C_P(H)B, H] = [B, H]$ . By assumption, we have  $G' \subseteq C_P(N)N$ . By Remark 3.6, this yields

$$[P, H] = [G', H] \subseteq [C_P(N)N, H] \subseteq [N, H][C_P(N), H],$$

since for  $c \in C_P(N)$ ,  $n \in N$  and  $h \in H$ , we have  $[cn, h] = c[n, h]c^{-1}[c, h] = [n, h][c, h]$ . Hence  $[P, H] \subseteq [B, H][C_P(N), H] \subseteq BC_P(N)$  follows, which yields

$$B = [C_P(H), [P, H]] \subseteq [C_P(H), BC_P(N)] = [C_P(H), B] \subseteq [P, B].$$

Hence  $B = 1$  follows, which yields  $P = C_P(H) * [P, H]$ . By Remark 3.6, this implies  $G = C_P(H) * H[P, H] = C_P(H) * O^p(G)$ . □

By Lemma 3.18, the properties given in Proposition 3.20 hold whenever  $\text{soc}(ZFG)$  is an ideal in  $FG$ . We conclude this section with a result on  $p$ -groups, which is an immediate consequence of Lemma 3.18:

**Lemma 3.21.** *If  $G$  is a  $p$ -group satisfying  $\text{soc}(ZFG) \trianglelefteq FG$ , then  $G$  is metabelian.*

*Proof.* Let  $A$  be a maximal abelian normal subgroup of  $G$ . Since  $C_G(A) = A$  holds, Lemma 3.18 yields  $G' \subseteq A$ . In particular,  $G'$  is abelian. □

**3.6. Special case  $G' \subseteq Z(P)$ .** Let  $G = P \rtimes H$  be a finite group with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$ . In this section, we show that  $\text{soc}(ZFG)$  is an ideal in  $FG$  if  $G' \subseteq Z(P)$  holds.

**Lemma 3.22.**

- (i) *Let  $g \in G$  with  $g_p \in Z(P)$ . Then  $[g] = [h] \cdot [g_p]$  holds for  $h \in H \cap gP$ .*
- (ii) *For  $u \in Z(P)$  and  $h \in C_G(H)$ , we have  $h[u] \subseteq [hu]$ .*
- (iii) *Assume  $[P, G] \subseteq Z(P)$ . Let  $h \in C_G(H)$  and write  $[h] = U_h h$  with  $U_h := \{[a, h] : a \in G\}$ . Then  $U_h$  is a normal subgroup of  $G$ .*

*Proof.*

- (i) By Remark 3.6,  $gP$  is a  $p'$ -section of  $G$ . In particular,  $[h]$  is the unique  $p'$ -conjugacy class contained in  $gP$  and hence  $[g_{p'}] = [h]$  follows. Since  $H$  is abelian, we have  $g_{p'} = uhu^{-1}$  for some  $u \in P$ . Due to  $g_p \in Z(P)$ , this yields  $g = uhg_pu^{-1}$  and hence  $[g] = [hg_p]$ . We may therefore assume  $g_{p'} = h$ . For  $x = p_x h_x$  with  $p_x \in P$  and  $h_x \in H$ , we have  $xgx^{-1} = p_x h p_x^{-1} \cdot h_x g_p h_x^{-1}$ . This yields

$$[g] = \{p_x h p_x^{-1} : p_x \in P\} \cdot \{h_x g_p h_x^{-1} : h_x \in H\} = [h] \cdot [g_p].$$

- (ii) Let  $u' \in [u]$ . Due to  $u \in Z(P)$ , there exists an element  $h' \in H$  with  $h'uh'^{-1} = u'$  (see Remark 3.6). Since  $h$  and  $h'$  commute, we obtain  $hu' = h'huh'^{-1} \in [hu]$ .
- (iii) We have  $U_h = \{[a, h] : a \in P\}$ . As  $[p_1 p_2, h] = [p_1, h] \cdot [p_2, h]$  holds for all  $p_1, p_2 \in P$ ,  $U_h$  is a subgroup of  $G'$ . Since the elements of  $P$  centralize  $U_h \subseteq [P, G] \subseteq Z(P)$  and conjugation with elements of  $H$  permutes the elements  $[a, h]$  with  $a \in P$ , it follows that  $U_h$  is normal in  $G$ . □

**Corollary 3.23.** *Let  $g \in G$  with  $g_p \in Z(P)$ . For  $y \in ZFG$  with  $y \cdot [g_p]^+ = 0$ , we have  $y \cdot [g]^+ = 0$ .*

*Proof.* The group  $P$  acts on  $[g]$  by conjugation with orbits of the form  $[g_{p'}]u$  with  $u \in P$  (see Lemma 3.22). In particular,  $[g]$  is a disjoint union of sets of this form. Hence  $y \cdot [g_p]^+$  implies  $y \cdot [g]^+ = 0$ .  $\square$

**Lemma 3.24.** *Let  $y = \sum_{g \in G} a_g g \in \text{soc}(ZFG)$ . For  $h \in C_G(H)$  and  $u \in Z(P)$ , we have  $a_{hu} = a_h$ .*

*Proof.* We may assume  $u \neq 1$ . By Remark 3.6,  $m := |[u]|$  is not divisible by  $p$ . Hence we have  $b_{[u^{-1}]} = [u^{-1}]^+ - m \cdot 1$  (see Theorem 3.14) and the coefficient of  $h$  in  $y \cdot b_{[u^{-1}]} = 0$  is given by

$$\sum_{u' \in [u]} a_{hu'} - ma_h = m(a_{hu} - a_h),$$

since the elements in  $h[u]$  are conjugate by Lemma 3.22(ii). Since  $p$  does not divide  $m$ , we obtain  $a_{hu} = a_h$ .  $\square$

**Theorem 3.25.** *If  $G = C_G(H)Z(P)$  holds, then  $\text{soc}(ZFG) \subseteq Z(P)^+ \cdot FG$  follows. In particular, if we have  $G' \subseteq Z(P)$ , then  $\text{soc}(ZFG)$  is an ideal in  $FG$ .*

*Proof.* Consider an element  $y = \sum_{g \in G} a_g g \in \text{soc}(ZFG)$ . Let  $g \in G$  and write  $g = cz$  with  $c \in C_G(H)$  and  $z \in Z(P)$ . By Lemma 3.24, we have  $a_g = a_{cz} = a_c$ . Hence  $y \in Z(P)^+ \cdot FG$  follows. If additionally  $G' \subseteq Z(P)$  holds, then  $\text{soc}(ZFG) \subseteq Z(P)^+ \cdot FG \subseteq (G')^+ \cdot FG$  follows, so  $\text{soc}(ZFG)$  is an ideal in  $FG$  (see Lemma 3.3).  $\square$

This proves the first part of Theorem C. The next example shows that the condition  $G' \subseteq Z(P)$  is not necessary for  $\text{soc}(ZFG) \trianglelefteq FG$ .

**Example 3.26.** Let  $F$  be an algebraically closed field of characteristic  $p = 3$  and consider the group  $G = \text{SmallGroup}(216, 86)$  in GAP [12]. We have  $G = G' \rtimes H$ , where  $G'$  is the extraspecial group of order 27 and exponent three, and  $H \cong C_8$  permutes the nontrivial elements of  $G'/G''$  transitively and acts on  $G'' = Z(G')$  by inversion. In particular,  $G'$  is nonabelian. For  $h \in H$ , we set  $S_h := \text{soc}(ZFG) \cap FhG'$ . Due to the  $H$ -grading of  $FG$  introduced in Remark 3.15, it suffices to show  $S_h = F(hG')^+$  for all  $h \in H$ . Clearly, we have  $(hG')^+ \in S_h$ . The derived subgroup  $G'$  decomposes into the  $G$ -conjugacy classes  $\{1\}$ ,  $G'' \setminus \{1\}$  and  $G' \setminus G''$ . For  $1 \neq h \in H$ , the coset  $hG'$  consists of a single conjugacy class for  $\text{ord}(h) = 8$  and of two conjugacy classes for  $\text{ord}(h) \in \{2, 4\}$ . In the first case, we directly obtain  $S_h = F(hG')^+$ . In the latter case, we have  $[h]^+ \cdot (G'')^+ = (hG')^+ \neq 0$ , which implies  $[h]^+ \notin \text{soc}(ZFG)$  since  $(G'')^+ \in J(ZFG)$  holds. Since  $(hG')^+ - [h]^+ \notin \text{soc}(ZFG)$  holds as well,  $S_h = F(hG')^+$  follows. Moreover, this shows  $(G'')^+ \notin \text{soc}(ZFG)$  and hence  $S_1 = F(G')^+$  follows as well. By Lemma 3.3,  $\text{soc}(ZFG)$  is an ideal of  $FG$ .

**3.7. Quotient groups.** Let  $G$  be a finite group of the form  $P \rtimes H$  with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$ . We fix a normal subgroup  $N \trianglelefteq G$  with quotient group  $\bar{G} := G/N$ . Our aim is to study the transition to the group algebra  $F\bar{G}$ . The image of an element  $g \in G$  in  $\bar{G}$  will be denoted by  $\bar{g}$  (similarly for subsets of  $G$ ). Note that  $\bar{G}$  is of the form  $\bar{P} \rtimes \bar{H}$  with  $\bar{P} \in \text{Syl}_p(\bar{G})$  and the abelian  $p'$ -group  $\bar{H}$ . In the following, we consider the canonical projection map

$$\nu_N: FG \rightarrow F\bar{G}, \quad \sum_{g \in G} a_g g \mapsto \sum_{g \in G} a_g \cdot gN,$$

together with its adjoint map  $\nu_N^*: F\bar{G} \rightarrow FG$ , which is defined by requiring  $\lambda(\nu_N^*(x)y) = \bar{\lambda}(x\nu_N(y))$  for all  $x \in F\bar{G}$  and  $y \in FG$ . Here,  $\lambda$  and  $\bar{\lambda}$  denote the symmetrizing linear forms of  $FG$  and  $F\bar{G}$  given in (2.1), respectively. It is easily verified that  $\nu_N^*$  is given by

$$\nu_N^*: F\bar{G} \rightarrow FG, \quad \sum_{gN \in \bar{G}} a_{gN} \cdot gN \mapsto \sum_{g \in G} a_{gN} \cdot g.$$

Note that  $\nu_N^*$  is a linear map with image  $N^+ \cdot FG$  and that it is injective as  $\nu_N$  is surjective.

**Remark 3.27.** For  $a \in F\bar{G}$ , it is easily seen that  $a \in (\bar{G}')^+ \cdot F\bar{G}$  is equivalent to  $\nu_N^*(a) \in (G')^+ \cdot FG$ .

If  $\text{soc}(ZFG)$  is an ideal in  $FG$ , then  $\text{Ann}_{ZF\bar{G}}(\nu_N(J(ZFG)))$  is an ideal in  $F\bar{G}$  by [3, Proposition 2.10]. For  $C \in \text{Cl}(G)$  with  $C \not\subseteq O_{p'}(G)$ , let  $b_C$  denote the associated element of  $J(ZFG)$  (see Definition 3.13) and consider the basis  $B := \{b_C : C \in \text{Cl}(G), C \not\subseteq O_{p'}(G)\}$  of  $J(ZFG)$  (see Theorem 3.14). Clearly,  $\nu_N(J(ZFG))$  is spanned by the images of the elements in  $B$ . We now derive a more convenient generating set.

**Lemma 3.28.** *Let  $C \in \text{Cl}(G)$  be a conjugacy class with  $C \not\subseteq O_{p'}(G)$ . We have  $b_C \notin \text{Ker}(\nu_N)$  if and only if  $\bar{C} \not\subseteq O_{p'}(\bar{G})$  holds and  $k := |C|/|\bar{C}|$  is not divisible by  $p$ . In this case, the basis element  $b_{\bar{C}}$  of  $J(ZF\bar{G})$  corresponding to  $\bar{C} \in \text{Cl}(\bar{G})$  is well-defined and we have  $\nu_N(b_C) = k \cdot b_{\bar{C}}$ .*

*Proof.* Observe that  $\bar{C}$  is indeed a conjugacy class of  $\bar{G}$  and that  $\nu_N(C^+) = k \cdot \bar{C}^+$  with  $k := |C|/|\bar{C}|$  holds. Suppose first that  $p$  divides  $|C|$ , so  $b_C = C^+$  holds. Then  $\nu_N(b_C) \neq 0$  is equivalent to  $k \not\equiv 0 \pmod{p}$ , and in this case we have  $|\bar{C}| \equiv 0 \pmod{p}$ . Since  $O_{p'}(\bar{G}) \subseteq Z(\bar{G})$  holds, this implies  $\bar{C} \not\subseteq O_{p'}(\bar{G})$ . Moreover, we have  $b_{\bar{C}} = \bar{C}^+$  and thus  $\nu_N(b_C) = k \cdot b_{\bar{C}}$ .

It remains to consider the case  $C \subseteq C_G(P)$ . There, we have  $\bar{C} \subseteq C_{\bar{G}}(\bar{P})$ . If  $\bar{C} \not\subseteq O_{p'}(\bar{G})$  holds, then  $b_{\bar{C}}$  is defined, and we have  $b_C = C^+ - |C| \cdot g_{p'}$  and  $b_{\bar{C}} = \bar{C}^+ - |\bar{C}| \cdot \bar{g}_{p'}$  for  $g \in C$ . This shows that  $\nu_N(b_C) = k \cdot b_{\bar{C}}$  holds. If, in addition,  $k \not\equiv 0 \pmod{p}$ , then  $\nu_N(b_C) \neq 0$  follows. Suppose conversely that  $\nu_N(b_C) \neq 0$  holds. We write  $C = g_{p'}D$  for  $g_{p'} \in O_{p'}(G)$  and  $D \in \text{Cl}(G)$  with  $D \subseteq Z(P)$  (see Remark 3.12). Assume that  $\bar{C} \subseteq O_{p'}(\bar{G})$  holds. Then we have  $\bar{D} = \bar{g}_{p'}^{-1}\bar{C} \subseteq O_{p'}(\bar{G})$  due to  $\bar{g}_{p'} \in O_{p'}(\bar{G})$ . As  $D$  consists of  $p$ -elements, we must have  $\bar{D} = \{1\}$ , which yields the contradiction  $\nu_N(b_C) = \nu_N(g_{p'}D^+ - |D| \cdot g_{p'}) = 0$ . This shows that  $\bar{C} \not\subseteq O_{p'}(\bar{G})$  holds. Hence we have  $\nu_N(b_C) = k \cdot b_{\bar{C}}$ , so that  $k \not\equiv 0 \pmod{p}$ .  $\square$

**Definition 3.29.** Set  $\text{Cl}_{p',N}(G) := \{C \in \text{Cl}(G) : C \not\subseteq O_{p'}(G) \text{ and } b_C \notin \text{Ker}(\nu_N)\}$  and let

$$\text{Cl}_{p',N}^+(G) := \{b_C : C \in \text{Cl}_{p',N}(G)\}$$

be the set of corresponding basis elements of  $J(ZFG)$  (see Definition 3.13). By  $\bar{\text{Cl}}_{p',N}(G) \subseteq \text{Cl}(\bar{G})$ , we denote the set of images of the conjugacy classes in  $\text{Cl}_{p',N}(G)$  and set

$$\bar{\text{Cl}}_{p',N}^+(G) := \{b_{\bar{C}} : \bar{C} \in \bar{\text{Cl}}_{p',N}(G)\},$$

where  $b_{\bar{C}}$  denotes the basis element of  $J(ZF\bar{G})$  corresponding to  $\bar{C}$ .

If  $N$  is a  $p$ -group, the  $p'$ -conjugacy classes of length divisible by  $p$  in  $\text{Cl}_{p',N}(G)$  can be easily characterized:

**Lemma 3.30.** *Consider a normal  $p$ -subgroup  $N$  of  $G$  and let  $C \not\subseteq C_G(P)$  be a  $p'$ -conjugacy class. Then we have  $C \in \text{Cl}_{p',N}(G)$  if and only if  $C \subseteq C_G(N)$  holds.*

*Proof.* If  $C \not\subseteq C_G(N)$  holds, we have  $\nu_N(b_C) = \nu_N(C^+) = 0$  by Lemma 3.16, so  $C \notin \text{Cl}_{p',N}(G)$ . Now let  $h \in C \subseteq C_G(N)$ . Since  $h$  is a  $p'$ -element, [5, Theorem 5.3.15] implies  $C_{G/N}(hN) = C_G(h)N/N = C_G(h)/N$  and hence  $|\bar{C}| = |G/N : C_{G/N}(hN)| = |G : C_G(h)| = |C|$ . Thus we have  $\nu_N(b_C) = \nu_N(C^+) = \bar{C}^+ \neq 0$ , which yields  $C \in \text{Cl}_{p',N}(G)$ .  $\square$

Now let  $N$  again be an arbitrary normal subgroup of  $G$ . We obtain the following necessary condition for  $\text{soc}(ZFG) \trianglelefteq FG$ :

**Theorem 3.31.** *We have*

$$\text{Ann}_{ZF\bar{G}}(\nu_N(J(ZFG))) = \text{Ann}_{ZF\bar{G}}(\overline{\text{Cl}}_{p',N}^+(G)) =: A.$$

If  $\text{soc}(ZFG)$  is an ideal of  $FG$ , we have  $A \subseteq (\bar{G}')^+ \cdot F\bar{G}$ .

*Proof.* Clearly, the elements  $\nu_N(b_C)$  with  $C \in \text{Cl}_{p',N}(G)$  span  $\nu_N(J(ZFG))$ . For  $C \in \text{Cl}_{p',N}(G)$  and  $y \in F\bar{G}$ , we have  $y \cdot \nu_N(b_C) = 0$  if and only if  $y \cdot b_{\bar{C}} = 0$  holds (see Lemma 3.28). This implies  $A = \text{Ann}_{ZF\bar{G}}(\nu_N(J(ZFG)))$ . Now assume that  $\text{soc}(ZFG)$  is an ideal in  $FG$ . By [3, Proposition 2.10],  $A$  is an ideal in  $F\bar{G}$ , so by [9, Lemma 2.1], we have  $K(F\bar{G}) \cdot A = 0$ . As in the proof of Lemma 3.3, this implies  $A \subseteq (\bar{G}')^+ \cdot F\bar{G}$ .  $\square$

As a first application, we give an alternative proof of the following special case of [3, Proposition 2.10]:

**Corollary 3.32.** *Let  $\text{soc}(ZFG)$  be an ideal of  $FG$ . Then  $\text{soc}(ZF\bar{G}) \trianglelefteq F\bar{G}$  holds.*

*Proof.* Since  $\overline{\text{Cl}}_{p',N}^+(G)$  is a subset of  $J(ZF\bar{G})$ , Theorem 3.31 yields

$$\text{soc}(ZF\bar{G}) = \text{Ann}_{ZF\bar{G}} J(ZF\bar{G}) \subseteq \text{Ann}_{ZF\bar{G}}(\overline{\text{Cl}}_{p',N}^+(G)) \subseteq (\bar{G}')^+ \cdot F\bar{G}$$

and we obtain  $\text{soc}(ZF\bar{G}) \trianglelefteq F\bar{G}$  by Lemma 3.3.  $\square$

**3.8. Central products.** Let  $G$  be a finite group. We consider the question when  $\text{soc}(ZFG)$  is an ideal of  $FG$  in case that  $G = G_1 * G_2$  is a central product of two subgroups  $G_1$  and  $G_2$ . Central products will play an important role throughout our investigation, for instance in the decomposition of  $G$  given in Theorem D.

**Theorem 3.33.** *Let  $G = G_1 * G_2$  be the central product of  $G_1$  and  $G_2$ . Then  $\text{soc}(ZFG) \trianglelefteq FG$  is equivalent to  $\text{soc}(ZFG_i) \trianglelefteq FG_i$  for  $i = 1, 2$ .*

*Proof.* First assume that  $\text{soc}(ZFG_i)$  is an ideal in  $FG_i$  for  $i = 1, 2$ . By [3, Proposition 1.9], this implies

$$\text{soc}(Z(FG_1 \otimes_F FG_2)) \trianglelefteq FG_1 \otimes_F FG_2.$$

Since  $F(G_1 \times G_2) \cong FG_1 \otimes_F FG_2$  holds, this yields  $\text{soc}(ZF(G_1 \times G_2)) \trianglelefteq F(G_1 \times G_2)$ . The group  $G$  is isomorphic to a quotient group of  $G_1 \times G_2$ , so  $\text{soc}(ZFG)$  is an ideal in  $FG$  by Corollary 3.32.

Now assume conversely that  $\text{soc}(ZFG)$  is an ideal of  $FG$ . By Corollary 3.5,  $G$  is of the form  $P \rtimes H$  with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$ . First suppose that  $O_{p'}(G) = 1$  holds. Then  $Z := G_1 \cap G_2 \subseteq Z(G) \subseteq C_G(P) = Z(P)$  is a  $p$ -group. We consider the canonical projection  $\nu := \nu_{G_2}: FG \rightarrow F[G/G_2]$ . By Theorem 3.31, we have

$$\text{Ann}_{ZF[G/G_2]}(\nu(J(ZFG))) \subseteq ([G/G_2]')^+ \cdot F[G/G_2]. \quad (3.1)$$

Note that there is a canonical isomorphism  $G_1/Z \cong G/G_2$ . Furthermore, we have  $ZFG_1 \subseteq ZFG$  and  $\nu(ZFG_1) = \nu(ZFG)$ , so also  $\nu(J(ZFG_1)) = \nu(J(ZFG))$  holds. Hence we have

$$\text{Ann}_{ZF[G_1/Z]}(\nu_1(J(ZFG_1))) \subseteq ([G_1/Z]')^+ \cdot F[G_1/Z],$$

where  $\nu_1: FG_1 \rightarrow F[G_1/Z]$  denotes the canonical projection. Let  $x_1 \in \text{soc}(ZFG_1)$  and observe that  $G'_1$  is a  $p$ -group. By Corollary 3.19, we have  $x_1 \in Z^+ \cdot FG_1 = \nu_1^*(F[G_1/Z])$ . Let  $y_1 \in F[G_1/Z]$  with  $x_1 = \nu_1^*(y_1)$ . Then [3, Remark 2.9] yields

$$y_1 \in \text{Ann}_{ZF[G_1/Z]}(\nu_1(J(ZFG_1))) \subseteq ([G_1/Z]')^+ \cdot F[G_1/Z].$$

By Remark 3.27, this yields  $x_1 \in (G'_1)^+ \cdot FG_1$  and hence  $\text{soc}(ZFG_1)$  is an ideal in  $FG_1$  (see Lemma 3.3). By symmetry, we obtain  $\text{soc}(ZFG_2) \trianglelefteq FG_2$ .

Now we consider the general case. For  $\bar{G} := G/O_{p'}(G)$ , we have  $\bar{G} = \bar{G}_1 * \bar{G}_2$  with  $\bar{G}_i := G_i O_{p'}(G)/O_{p'}(G)$  ( $i = 1, 2$ ). Note that  $\bar{G}_i \cong G_i/O_{p'}(G) \cap G_i \cong G_i/O_{p'}(G_i)$  follows since  $O_{p'}(G) \cap G_i = O_{p'}(G_i)$  holds. By the above, we obtain  $\text{soc}(ZF\bar{G}_i) \trianglelefteq F\bar{G}_i$ . Since  $G'$  is a  $p$ -group, also  $G'_1$  and  $G'_2$  are  $p$ -groups. Lemma 3.10 then yields  $\text{soc}(ZFG_i) \trianglelefteq FG_i$  for  $i = 1, 2$ . □

**Remark 3.34.** For  $G \cong G_1 \times G_2$ , the statement of Theorem 3.33 is a special case of [3, Proposition 1.9].

#### 4. GROUPS OF PRIME POWER ORDER

Let  $F$  be an algebraically closed field of characteristic  $p > 0$ . In this section, we classify the finite  $p$ -groups  $G$  for which  $\text{soc}(ZFG)$  is an ideal in  $FG$  (see Theorem B). Additionally, these results will be generalized to arbitrary finite groups (see Theorem C). First we prove that the property  $\text{soc}(ZFG) \trianglelefteq FG$  is preserved under isoclinism (see Section 4.1). Subsequently, we distinguish the cases  $p \geq 3$  (see Section 4.2) and  $p = 2$  (see Section 4.3).

**4.1. Isoclinism.** Let  $G$  be a finite  $p$ -group. The aim of this section is to show that the property  $\text{soc}(ZFG) \trianglelefteq FG$  is invariant under isoclinism in the following sense: If  $Q$  is a finite  $p$ -group isoclinic to  $G$ , then  $\text{soc}(ZFQ) \trianglelefteq FQ$  holds precisely if we have  $\text{soc}(ZFG) \trianglelefteq FG$ . The proof of this statement is based on some observations on the center of  $G$  and the transition to the quotient group  $\bar{G} := G/Z(G)$ .

**Lemma 4.1.**

- (i) We have  $\text{soc}(ZFG) \subseteq Z(G)^+ \cdot FG$ .
- (ii)  $\text{soc}(ZFG)$  is an ideal of  $FG$  if and only if  $\text{soc}(ZFG) = (Z(G)G')^+ \cdot FG$  holds.

*Proof.* The first statement follows by Corollary 3.19. Now let  $\text{soc}(ZFG)$  be an ideal of  $FG$ . Lemma 3.3 then yields  $\text{soc}(ZFG) \subseteq (G')^+ \cdot FG$ . Together with (i), this implies  $\text{soc}(ZFG) \subseteq (Z(G)G')^+ \cdot FG$ , and by Corollary 3.19, we obtain equality. Conversely,  $(Z(G)G')^+ \cdot FG$  is obviously an ideal in  $FG$ . □

In the given situation, we have

$$\text{Cl}_{p'} := \text{Cl}_{p',Z(G)}(G) = \left\{ C \in \text{Cl}(G) : C \not\subseteq Z(G), |C| = |\bar{C}| \right\}.$$

Note that the length of every conjugacy class in  $\text{Cl}_{p'}$  is a nontrivial power of  $p$ . Let  $\bar{\text{Cl}}_{p'} := \bar{\text{Cl}}_{p',Z(G)}(G)$  be the set of images of the classes in  $\text{Cl}_{p'}$  and denote by  $\bar{\text{Cl}}_{p'}^+ :=$

$\overline{\text{Cl}}_{p',Z(G)}^+(G)$  the corresponding class sums in  $F\bar{G}$ . In this situation, the implication given in Theorem 3.31 is an equivalence:

**Lemma 4.2.** *The socle  $\text{soc}(ZFG)$  is an ideal in  $FG$  if and only if  $\text{Ann}_{ZF\bar{G}}(\overline{\text{Cl}}_{p'}^+) \subseteq (\bar{G}')^+ \cdot F\bar{G}$  holds.*

*Proof.* Consider the map  $\nu_{Z(G)}^*: F\bar{G} \rightarrow FG$  introduced in Section 3.7. Lemma 4.1 yields

$$\text{soc}(ZFG) \subseteq Z(G)^+ \cdot FG = \text{Im } \nu_{Z(G)}^*.$$

By [3, Remark 2.9], we therefore obtain

$$\text{soc}(ZFG) = \nu_{Z(G)}^* \left( \text{Ann}_{ZF\bar{G}}(\overline{\text{Cl}}_{p'}^+) \right).$$

By Remark 3.27, we have  $\text{Ann}_{ZF\bar{G}}(\overline{\text{Cl}}_{p'}^+) \subseteq (\bar{G}')^+ \cdot F\bar{G}$  if and only if  $\text{soc}(ZFG) \subseteq (G')^+ \cdot FG$  holds, which is equivalent to  $\text{soc}(ZFG) \trianglelefteq FG$  by Lemma 3.3.  $\square$

Now we proceed to the main result of this section. Recall that two finite  $p$ -groups  $G_1$  and  $G_2$  are isoclinic if there exist isomorphisms  $\varphi: G'_1 \rightarrow G'_2$  and  $\beta: G_1/Z(G_1) \rightarrow G_2/Z(G_2)$  such that whenever  $\beta(a_1Z(G_1)) = a_2Z(G_2)$  and  $\beta(b_1Z(G_1)) = b_2Z(G_2)$  hold for  $a_1, b_1 \in G_1$  and  $a_2, b_2 \in G_2$ , then  $\varphi([a_1, b_1]) = [a_2, b_2]$  follows. We set  $\bar{G}_i := G_i/Z(G_i)$  and write  $\text{Cl}_{p',i}$  and  $\overline{\text{Cl}}_{p',i}$  to distinguish the sets  $\text{Cl}_{p'}$  and  $\overline{\text{Cl}}_{p'}$  for  $i \in \{1, 2\}$ .

**Theorem 4.3.** *Let  $G_1$  and  $G_2$  be finite isoclinic  $p$ -groups. Then  $\text{soc}(ZFG_1) \trianglelefteq FG_1$  is equivalent to  $\text{soc}(ZFG_2) \trianglelefteq FG_2$ .*

*Proof.* Let  $\varphi: G'_1 \rightarrow G'_2$  and  $\beta: \bar{G}_1 \rightarrow \bar{G}_2$  be the corresponding isomorphisms. We first show that  $\overline{\text{Cl}}_{p',1}$  and  $\overline{\text{Cl}}_{p',2}$  are in bijective correspondence under  $\beta$ . Let  $C_1 \in \text{Cl}_{p',1}$  and set  $\bar{C}_1$  to be its image in  $\bar{G}_1$ . Then  $\bar{C}_2 := \beta(\bar{C}_1)$  is a conjugacy class of  $\bar{G}_2$ . Consider a preimage  $C_2 \in \text{Cl}(G_2)$  of  $\bar{C}_2$ . Let  $x_2 \in C_2$  and assume that  $1 \neq [x_2, g_2] \in Z(G_2)$  holds for some  $g_2 \in G_2$ . Choose elements  $x_1 \in C_1$  and  $g_1 \in G_1$  with  $\beta(x_1Z(G_1)) = x_2Z(G_2)$  and  $\beta(g_1Z(G_1)) = g_2Z(G_2)$ . We then obtain  $\varphi([x_1, g_1]) = [x_2, g_2] \in Z(G_2) \setminus \{1\}$ . Note that  $\beta([x_1, g_1]Z(G_1)) = [x_2, g_2]Z(G_2) = Z(G_2)$  holds, so we have  $1 \neq [x_1, g_1] \in Z(G_1)$ . This implies  $|\bar{C}_1| < |C_1|$ , which is a contradiction to  $C_1 \in \text{Cl}_{p',1}$ . Hence we obtain  $\bar{C}_2 \in \overline{\text{Cl}}_{p',2}$ . The other implication follows by symmetry.

Extending  $\beta$   $F$ -linearly gives rise to an  $F$ -algebra isomorphism  $\hat{\beta}: F\bar{G}_1 \rightarrow F\bar{G}_2$ . By the above, we have  $\hat{\beta}(\overline{\text{Cl}}_{p',1}^+) = \overline{\text{Cl}}_{p',2}^+$ . Now if  $\text{soc}(ZFG_1)$  is an ideal of  $FG_1$ , Lemma 4.2 implies  $\text{Ann}_{ZF\bar{G}_1}(\overline{\text{Cl}}_{p',1}^+) \subseteq (\bar{G}'_1)^+ \cdot F\bar{G}_1$ . Applying the isomorphism  $\hat{\beta}$  yields  $\text{Ann}_{ZF\bar{G}_2}(\overline{\text{Cl}}_{p',2}^+) \subseteq (\bar{G}'_2)^+ \cdot F\bar{G}_2$ . By Lemma 4.2,  $\text{soc}(ZFG_2)$  is an ideal in  $FG_2$ . The other implication follows by symmetry.  $\square$

**4.2. Odd characteristic.** In this section, we assume that  $F$  is an algebraically closed field of odd characteristic  $p$ .

**Remark 4.4.** For an abelian  $p$ -group  $G$ , we have  $\prod_{g \in G} g = 1$  since every nontrivial element in  $G$  differs from its inverse and their product is the identity element.

**Proposition 4.5.** *Let  $G$  be a finite  $p$ -group of nilpotency class exactly two. Then there exists an element  $y \in ZFG$  with  $y \notin (G')^+ \cdot FG$  such that  $y \cdot S^+ = 0$  holds for all subgroups  $1 \neq S \subseteq G'$ .*



*Proof.* Since  $G'$  is a nontrivial  $p$ -group, there exists a nontrivial group homomorphism  $\alpha: G' \rightarrow F$ . We define an element  $y := \sum_{g \in G'} a_g g \in FG$  by setting  $a_g := \alpha(g)$  for  $g \in G'$  and  $a_g = 0$  otherwise. We have  $y \in FG' \subseteq FZ(G) \subseteq ZFG$ . Now consider a subgroup  $1 \neq S \subseteq G'$ . The coefficient of  $w \in G$  in the product  $y \cdot S^+$  is given by  $\sum_{s \in S} a_{ws^{-1}}$ . For  $w \notin G'$ , all summands are zero. For  $w \in G'$ , we obtain

$$\sum_{s \in S} a_{ws^{-1}} = \sum_{s \in S} \alpha(ws^{-1}) = |S| \cdot \alpha(w) + \sum_{s \in S} \alpha(s^{-1}) = \alpha\left(\prod_{s \in S} s^{-1}\right) = \alpha(1) = 0.$$

In the second and third step, we use that  $\alpha$  is a group homomorphism. The fourth equality is due to Remark 4.4. This implies  $y \cdot S^+ = 0$  as claimed.  $\square$

In this special situation, the condition given in Theorem 3.25 is in fact equivalent to  $\text{soc}(ZFG) \trianglelefteq FG$ :

**Theorem 4.6.** *Let  $G$  be a finite  $p$ -group. Then  $\text{soc}(ZFG)$  is an ideal in  $FG$  if and only if  $G$  has nilpotency class at most two.*

*Proof.* If  $G$  is of nilpotency class at most two, we have  $G' \subseteq Z(G)$  and hence  $\text{soc}(ZFG)$  is an ideal in  $FG$  by Theorem 3.25. For the converse implication, we use induction on the nilpotency class of  $G$ . Note that  $\bar{G} := G/Z(G)$  has nilpotency class  $c(G) - 1$ . First assume  $c(G) = 3$ . We apply Proposition 4.5 to the group  $\bar{G}$  and consider the element  $y \in ZF\bar{G}$  constructed therein. Let  $\bar{C} \in \overline{\text{Cl}}_{p'}'$  be a conjugacy class and let  $c \in \bar{C}$ . Since  $\bar{G}' \subseteq Z(\bar{G})$  holds, the map  $\gamma: \bar{G} \rightarrow \bar{G}$ ,  $g \mapsto [g, c]$  is a group homomorphism and hence we have

$$\bar{C} = \{g c g^{-1} : g \in \bar{G}\} = \{[g, c]c : g \in \bar{G}\} = S c,$$

where  $S := \text{Im } \gamma$  is a subgroup of  $\bar{G}'$ . Note that we have  $|S| = |\bar{C}| > 1$ . By Proposition 4.5, we have  $y \cdot S^+ = 0$  and hence  $y \cdot \bar{C}^+ = y \cdot (S c)^+ = 0$ . Since  $y \notin (\bar{G}')^+ \cdot F\bar{G}$  holds,  $\text{soc}(ZFG)$  is not an ideal of  $FG$  (see Lemma 4.2). If  $G$  is of nilpotency class  $c(G) > 3$ , we obtain  $\text{soc}(ZF\bar{G}) \not\trianglelefteq F\bar{G}$  by induction. Corollary 3.32 then yields  $\text{soc}(ZFG) \not\trianglelefteq FG$ .  $\square$

**Remark 4.7.** The analogous construction fails for  $p = 2$  since the statement of Remark 4.4 does not hold for groups of even order.

**4.3. Characteristic  $p = 2$ .** Throughout, let  $F$  be an algebraically closed field of characteristic two. Unless otherwise stated, we assume that  $G$  is a finite 2-group.

**Remark 4.8.** Let  $C = \{f, g\}$  be a conjugacy class of length two of  $G$ . An inner automorphism of  $G$  either fixes both  $f$  and  $g$ , or it interchanges the two elements. For  $c := g f^{-1} \in G'$ , this yields  $C_G(f) = C_G(g) \subseteq C_G(c)$ . For  $h \in G \setminus C_G(f)$ , we have  $h c h^{-1} = h g f^{-1} h^{-1} = f g^{-1} = c^{-1}$ . This shows that the subgroup  $\langle c \rangle \subseteq G'$  is normal in  $G$ .

For every conjugacy class  $C := \{f, g\}$  of length two, we set  $Y_C := \langle g f^{-1} \rangle$ . In the following, we consider the subgroup

$$Y(G) := \langle Y_C : C \in \text{Cl}(G), |C| = 2 \rangle.$$

Note that  $Y(G)$  is characteristic in  $G$ . More precisely, we obtain the following:

**Lemma 4.9.** *We have  $Y(G) \subseteq Z(\Phi(G))$ . In particular,  $Y(G)$  is abelian.*

*Proof.* Note that  $Y(G) \subseteq G' \subseteq \Phi(G)$  holds. Now let  $C = \{f, g\}$  be a conjugacy class of length two. Since  $C_G(f)$  is a maximal subgroup of  $G$ , Remark 4.8 yields  $\Phi(G) \subseteq C_G(f) \subseteq C_G(g f^{-1})$  and hence  $\Phi(G)$  centralizes  $Y_C$ . Thus  $Y(G)$  is contained in the center of  $\Phi(G)$ , so in particular, it is abelian.  $\square$

**Lemma 4.10.** *We have  $\text{soc}(ZFG) \subseteq Y(G)^+ \cdot FG$ .*

*Proof.* Let  $y = \sum_{g \in G} a_g g \in \text{soc}(ZFG)$ . For a conjugacy class  $C = \{f, g\}$  of length two, we have  $c := gf^{-1} \in Y(G)$  and the condition  $y \cdot C^+ = 0$  yields  $a_x = a_{xc^{-1}}$  for all  $x \in G$ . By induction, this implies  $a_x = a_{xc_1^{-1} \dots c_n^{-1}}$  for every  $x \in G$  and all elements  $c_1, \dots, c_n$  arising from  $G$ -conjugacy classes of length two as above. This shows that  $y$  has constant coefficients on the cosets of  $Y(G)$ , that is, we obtain  $y \in Y(G)^+ \cdot FG$ .  $\square$

With this preliminary result, we obtain the following characterization of the 2-groups  $G$  for which  $\text{soc}(ZFG)$  is an ideal in  $FG$ .

**Theorem 4.11.** *The socle  $\text{soc}(ZFG)$  is an ideal in  $FG$  if and only if  $G' \subseteq Y(G)Z(G)$  holds.*

*Proof.* Suppose  $G' \not\subseteq Y(G)Z(G)$ , so  $Y(G)Z(G) \cap G'$  is a proper subgroup of  $G'$ . By [6, Theorem III.7.2], there exists a subgroup  $N \trianglelefteq G$  with  $Y(G)Z(G) \cap G' \subseteq N \subseteq G'$  and  $|G' : N| = 2$ . We set  $M := Y(G)Z(G)N$ . Note that  $M^+ \in ZFG$  holds since  $M$  is a normal subgroup of  $G$ . We now show that  $M^+$  annihilates the basis of  $J(ZFG)$  given in Theorem 3.14.

For  $z \in Z(G) \subseteq M$ , we have  $(1 + z) \cdot M^+ = 0$ . For a  $G$ -conjugacy class  $C = \{f, g\}$  of length two, we obtain  $C^+ \cdot Y(G)^+ = fY(G)^+ + gY(G)^+ = 0$  since  $gf^{-1} \in Y(G)$  holds. Hence  $M^+$  annihilates  $C^+$ . Every conjugacy class  $C \in \text{Cl}(G)$  with  $|C| \geq 4$  contains an even number of elements in every coset of  $N$  since  $C$  is contained in a coset of  $G'$  and  $|G' : N| = 2$  holds. This implies that  $C^+$  is annihilated by  $N^+$  and hence by  $M^+$ . Summarizing, we obtain  $M^+ \in \text{soc}(ZFG)$ . Moreover,  $M \cap G' = N \subsetneq G'$  implies  $M^+ \notin (G')^+ \cdot FG$ . By Lemma 3.3, this yields  $\text{soc}(ZFG) \not\subseteq FG$ .

Conversely, assume that  $G' \subseteq Y(G)Z(G)$  holds. By Lemmas 4.1 and 4.10, we have

$$\text{soc}(ZFG) \subseteq (Y(G)Z(G))^+ \cdot FG \subseteq (G')^+ \cdot FG$$

and hence  $\text{soc}(ZFG)$  is an ideal of  $FG$  (see Lemma 3.3).  $\square$

This completes the proof of Theorem B.

**Remark 4.12.** Similarly to the case of odd characteristic,  $\text{soc}(ZFG) \trianglelefteq FG$  holds if  $G$  has nilpotency class at most two.

However, the next example demonstrates that in contrast to the case of odd characteristic, the nilpotency class of a finite 2-group  $G$  for which  $\text{soc}(ZFG)$  is an ideal in  $FG$  can be arbitrarily large.

**Example 4.13.**

- (i) Let  $G = D_{2^n} = \langle r, s : r^{2^{n-1}} = s^2 = 1, srs = r^{-1} \rangle$  with  $n \in \mathbb{N}$  be the dihedral group of order  $2^n$ . For  $n \leq 2$ ,  $G$  is abelian and hence  $\text{soc}(ZFG) \trianglelefteq FG$  holds. For  $n \geq 3$ , we have  $G' = \langle r^2 \rangle = Y(G)Z(G)$  and hence  $\text{soc}(ZFG) \trianglelefteq FG$  follows by Theorem 4.11. The 2-groups of maximal class of a fixed order are isoclinic. Therefore, by Theorem 4.3,  $\text{soc}(ZFG)$  is an ideal in  $FG$  if  $G$  is a semihedral or generalized quaternion 2-group.
- (ii) By [3, Theorem 4.12], every 2-group  $G$  of order at most 16 satisfies  $\text{soc}(ZFG) \trianglelefteq FG$ . Up to isomorphism, there exist 51 groups of order 32. Out of those, 7 groups are abelian and 26 groups have nilpotency class precisely two. Additionally, 13 groups satisfy the property  $G' \subseteq Y(G)Z(G)$ .

- (iii) Consider the holomorph  $G = \mathbb{Z}/8\mathbb{Z} \rtimes (\mathbb{Z}/8\mathbb{Z})^\times$  of  $\mathbb{Z}/8\mathbb{Z}$ , which has order 32. It has 11 conjugacy classes and we have  $|Z(G)| = 2$ . Since  $G/Z(G) \cong D_8 \times C_2$  has precisely 10 conjugacy classes, the images of the non-central conjugacy classes of  $G$  in  $G/Z(G)$  are pairwise distinct. For every such conjugacy class  $C$ , we therefore have  $Z(G)C \subseteq C$  and hence  $\nu_{Z(G)}(C^+) = 0$ . This proves  $J(ZFG)^2 = 0$ , so  $J(ZFG) = \text{soc}(ZFG)$  follows. In particular, we obtain  $\dim \text{soc}(ZFG) = \dim J(ZFG) = 10$ . Due to  $|G'| = 4$ , the space  $(G')^+ \cdot FG$  is eight-dimensional, so it does not contain  $\text{soc}(ZFG)$ . By Lemma 3.3,  $\text{soc}(ZFG)$  is not an ideal in  $FG$ .

We conclude this part with a generalization of Theorem 4.14 to arbitrary finite groups, which is a stronger variant of Theorem 3.25:

**Theorem 4.14.** *Let  $G$  be an arbitrary finite group which satisfies  $G' \subseteq Y(O_2(G))Z(O_2(G))$ . Then  $\text{soc}(ZFG)$  is an ideal of  $FG$ .*

*Proof.* The given condition implies  $G' \subseteq O_2(G)$ , so by Theorem 3.4, we have  $G = P \rtimes H$  with  $P := O_2(G) \in \text{Syl}_2(G)$  and an abelian 2'-group  $H$ . Note that  $G'$  is abelian as  $Y(P)$  is abelian (see Lemma 4.9). By Remark 3.6, we have

$$P = C_P(H)G' = C_P(H)Y(P)Z(P). \tag{4.1}$$

Since  $G'$  is abelian,  $C_P(H)'$  is normal in  $P$ . We consider the group  $\bar{P} := P/C_P(H)'$  and denote the image of  $S \subseteq P$  in  $\bar{P}$  by  $\bar{S}$ . Then we have

$$\overline{Y(P)} \subseteq \bar{P}' = \overline{(C_P(H)Y(P))'} = [\overline{C_P(H)}, \overline{Y(P)}] \subseteq [\bar{P}, \overline{Y(P)}].$$

This implies  $\overline{Y(P)} = 1$ , so  $Y(P) \subseteq C_P(H)'$  follows.

By (4.1), we then have  $[P, H] = [Z(P), H]$  and hence  $P = C_P(H)[Z(P), H]$  follows. Since  $C_P(H)$  centralizes  $W := H[Z(P), H] \subseteq HZ(P)$  and  $C_P(H) \cap [Z(P), H] = 1$  follows by [5, Theorem 5.3.6], we obtain  $G = C_P(H) \times W$ . It is then easily verified that  $Y(P) = Y(C_P(H))$  holds. With Dedekind's identity, we obtain

$$C_P(H)' \subseteq G' \cap C_P(H) \subseteq Y(C_P(H))Z(P) \cap C_P(H) \subseteq Y(C_P(H)) \cdot Z(C_P(H)).$$

By Theorem 4.11,  $\text{soc}(ZFC_P(H))$  is an ideal of  $FC_P(H)$ . Since  $\text{soc}(ZFW) \trianglelefteq FW$  follows by Theorem 3.25,  $\text{soc}(ZFG)$  is an ideal in  $FG$  by Theorem 3.33. □

This completes the proof of Theorem C.

### 5. DECOMPOSITION OF $G$ INTO A CENTRAL PRODUCT

Let  $F$  be an algebraically closed field of characteristic  $p > 0$ . We consider an arbitrary finite group  $G$  for which  $\text{soc}(ZFG)$  is an ideal in  $FG$ . By Theorem 3.4, we may write  $G = P \rtimes H$  with  $P \in \text{Syl}_p(G)$  and an abelian  $p'$ -group  $H$ . In this section, we prove Theorem D. Combined with the results on  $p$ -groups from the last section, it reduces our investigation to the case that  $G'$  is a Sylow  $p$ -subgroup of  $G$ .

**Theorem 5.1** (Theorem D). *We have  $G = C_P(H) * O^p(G)$ . Moreover,  $\text{soc}(ZFC_P(H))$  and  $\text{soc}(ZFO^p(G))$  are ideals in  $FC_P(H)$  and  $FO^p(G)$ , respectively. The socle of  $ZFG$  is explicitly given by*

$$\text{soc}(ZFG) = (Z(P)G')^+ \cdot FG.$$

*Proof.* By Proposition 3.20, we have  $G = C_P(H) * O^p(G)$ . Theorem 3.33 then implies that  $\text{soc}(ZFC_P(H))$  and  $\text{soc}(ZFO^p(G))$  are ideals in  $FC_P(H)$  and  $FO^p(G)$ , respectively. It therefore remains to determine the structure of  $\text{soc}(ZFG)$ . By the above decomposition, we have  $Z(C_P(H)) \subseteq Z(G)$ . By Corollary 3.19, we obtain  $\text{soc}(ZFG) \subseteq Z(C_P(H))^+ \cdot FG$ . Together with Lemma 3.3, this implies

$$\text{soc}(ZFG) \subseteq (Z(C_P(H))G')^+ \cdot FG \subseteq (Z(P)G')^+ \cdot FG.$$

In the last step, we used  $Z(P) = Z(C_P(H))Z([G', H]) \subseteq Z(C_P(H))G'$ . On the other hand, we have  $(Z(P)G')^+ \cdot FG \subseteq \text{soc}(ZFG)$  by Lemma 3.16, which completes the proof of Theorem D.  $\square$

This result on the structure of  $\text{soc}(ZFG)$  generalizes the corresponding statement in Lemma 4.1. Note that, by Theorem D, the hypothesis that  $\text{soc}(ZFG)$  is an ideal in  $FG$  implies  $\dim \text{soc}(ZFG) = |G : G'Z(G)|$ . In particular, the dimension of  $\text{soc}(ZFG)$  is a divisor of  $|G|$ . We also observe that in this situation,  $\text{soc}(ZFG)$  is a principal ideal of  $FG$  generated by a central element. Furthermore, we obtain the following reduction:

**Remark 5.2.** Since the structure of the  $p$ -group  $C_P(H)$  is determined by Theorem B, it suffices to investigate the group  $O^p(G)$ . Inductively, we may assume  $O^p(G) = G$ . By Remark 3.6, this yields  $P = G' = [G', H]$ . In particular,  $C_{G'}(H) \subseteq G'' \subseteq Z(G')$  follows (see [5, Theorem 5.2.3]), which implies  $C_{G'}(H) \subseteq Z(G)$ . If additionally  $O_{p'}(G) = 1$  holds, we obtain  $C_{G'}(H) = Z(G)$ .

Moreover, we state the following consequence of Theorem D:

**Theorem 5.3.** *We have  $\text{soc}(ZFP) \trianglelefteq FP$ . In particular, the group  $P$  is metabelian and its nilpotency class is at most two if  $p$  is odd.*

*Proof.* By Theorem D, we have  $P = C_P(H) * [P, H]$  and  $\text{soc}(ZFC_P(H))$  is an ideal in  $FC_P(H)$ . Since  $[P, H] \subseteq G'$  has nilpotency class at most two (see Proposition 3.20), we obtain  $\text{soc}(ZF[P, H]) \trianglelefteq F[P, H]$  by Theorem B. By Theorem 3.33, this yields  $\text{soc}(ZFP) \trianglelefteq FP$ . In particular, it follows that  $P$  is metabelian and that the nilpotency class of  $P$  is at most two if  $p$  is odd (see Theorem B).  $\square$

#### ACKNOWLEDGMENTS

The results of this paper form a part of the PhD thesis of the first author [2], which was supervised by the second author.

#### REFERENCES

- [1] Richard Brauer, *Some applications of the theory of blocks of characters of finite groups. I*, J. Algebra **1** (1964), no. 2, 152–167.
- [2] Sofia Brenner, *The socle of the center of a group algebra*, Dissertation, Friedrich-Schiller-Universität Jena, Deutschland, 2022.
- [3] Sofia Brenner and Burkhard Külshammer, *Ideals in the center of symmetric algebras*, Int. Electron. J. Algebra **34** (2023), 126–151.
- [4] Robert J. Clarke, *On the Radical of the Centre of a Group Algebra*, J. Lond. Math. Soc., II. Ser. **2** (1969), no. 1, 565–572.
- [5] Daniel Gorenstein, *Finite Groups*, Harper's Series in Modern Mathematics, Harper & Row, 1968.
- [6] Bertram Huppert, *Endliche Gruppen. I*, Grundlehren der Mathematischen Wissenschaften, vol. 134, Springer, 1967.
- [7] Shigeo Koshitani, *A Note on the Radical of the Centre of a Group Algebra*, J. Lond. Math. Soc. **18** (1978), no. 2, 243–246.

- [8] Burkhard Külshammer, *Group-theoretical descriptions of ring-theoretical invariants of group algebras*, Representation Theory of Finite Groups and Finite-Dimensional Algebras (Gerhard O. Michler and Claus M. Ringel, eds.), Progress in Mathematics, vol. 95, Birkhäuser, 1991, pp. 425–442.
- [9] ———, *Centers and radicals of group algebras and blocks*, Arch. Math. **114** (2020), 619–629.
- [10] Markus Linckelmann, *The Block Theory of Finite Group Algebras. Volume 1*, London Mathematical Society Student Texts, vol. 91, Cambridge University Press, 2018.
- [11] Donald S. Passman, *The Algebraic Structure of Group Rings*, John Wiley & Sons, 1977.
- [12] The GAP Group, *Gap – Groups, Algorithms, and Programming, Version 4.10.0*, 2018, <https://www.gap-system.org>.

— SOFIA BRENNER —

DEPARTMENT OF MATHEMATICS, TU DARMSTADT, GERMANY

*E-mail address:* [sofia.brenner@tu-darmstadt.de](mailto:sofia.brenner@tu-darmstadt.de)

INSTITUTE FOR MATHEMATICS, FRIEDRICH SCHILLER UNIVERSITY JENA, GERMANY

*E-mail address:* [sofia.bettina.brenner@uni-jena.de](mailto:sofia.bettina.brenner@uni-jena.de)

— BURKHARD KÜLSHAMMER —

INSTITUTE FOR MATHEMATICS, FRIEDRICH SCHILLER UNIVERSITY JENA, GERMANY

*E-mail address:* [kuelshammer@uni-jena.de](mailto:kuelshammer@uni-jena.de)