



# Annals of Representation Theory

JONATHAN BRUNDAN, WEIQIANG WANG & BENJAMIN WEBSTER 

## The nil-Brauer category

Volume 1, issue 1 (2024), p. 21-58

<https://doi.org/10.5802/art.2>

Communicated by Kevin Coulembier.

© The authors, 2024



This article is licensed under the  
CREATIVE COMMONS ATTRIBUTION (CC-BY) 4.0 LICENSE.  
<http://creativecommons.org/licenses/by/4.0/>



NTNU

*Annals of Representation Theory* is published by the  
Norwegian University of Science and Technology  
and a member of the  
Centre Mersenne for Open Scientific Publishing

e-ISSN: pending





# The nil-Brauer category

Jonathan Brundan, Weiqiang Wang\* and Benjamin Webster 

**ABSTRACT.** We introduce the nil-Brauer category and prove a basis theorem for its morphism spaces. This basis theorem is an essential ingredient required to prove that nil-Brauer categorifies the split  $\iota$ -quantum group of rank one. As this  $\iota$ -quantum group is a basic building block for  $\iota$ -quantum groups of higher rank, we expect that the nil-Brauer category will play a central role in future developments related to the categorification of quantum symmetric pairs.

## 1. INTRODUCTION

Throughout the article, we work over an integral domain  $\mathbb{k}$  in which 2 is invertible, and all categories, functors, etc. will be assumed to be  $\mathbb{k}$ -linear without further mention. The aim of the article is to introduce a new strict graded monoidal category, the *nil-Brauer category*, denoted  $\mathcal{NB}_t$  for a parameter  $t \in \mathbb{k}$ . It turns out that  $\mathcal{NB}_t$  is non-trivial only for  $t = 0$  or  $t = 1$ ; we assume this is the case for the remainder of the introduction.

The importance of the nil-Brauer category stems from results established in our subsequent work [11] relating  $\mathcal{NB}_t$  to  $U_q^{\iota}(\mathfrak{sl}_2)$ , the split  $\iota$ -quantum group of rank one corresponding to the symmetric pair  $(\mathrm{SL}_2, \mathrm{SO}_2)$ . Roughly speaking, the monoidal categories  $\mathcal{NB}_t$  ( $t = 0, 1$ ) play the same role for this, the most basic of all  $\iota$ -quantum groups, as the strict graded 2-category  $\mathfrak{U}(\mathfrak{sl}_2)$  introduced in [20, 16] plays for the ordinary quantum group  $U_q(\mathfrak{sl}_2)$ . To make a more precise statement, let  ${}_{\mathbb{Z}}\mathbf{U}_t^{\iota}$  be the modified  $\mathbb{Z}[q, q^{-1}]$ -form of  $U_q^{\iota}(\mathfrak{sl}_2)$  associated to weights of parity  $t \in \{0, 1\}$  arising as a special case of the constructions in [2, 3]. The algebra  $U_q^{\iota}(\mathfrak{sl}_2)$  is simply a polynomial algebra in one variable  $B$ , but

---

Manuscript received 2023-05-18, revised 2023-06-27 and accepted 2023-09-15.

*Keywords.* Brauer category, string calculus, categorification.

*2020 Mathematics Subject Classification.* 17B10, 17B37.

J.B. supported in part by NSF grant DMS-2101783. W.W. supported in part by the NSF grant DMS-2001351. B.W. supported by Discovery Grant RGPIN-2018-03974 from the Natural Sciences and Engineering Research Council of Canada. This research was also supported by Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development and by the Province of Ontario through the Ministry of Research and Innovation.

\* Corresponding author.

the integral form  ${}_{\mathbb{Z}}\mathbf{U}_t^{\iota}$  is not at all obvious; it has a basis as a free  $\mathbb{Z}[q, q^{-1}]$ -module given by the  $\iota$ -canonical basis which was computed explicitly in [4]. We will show in [11] that  $\mathcal{N}\mathcal{B}_t$  categorifies  ${}_{\mathbb{Z}}\mathbf{U}_t^{\iota}$  in the sense that the split Grothendieck ring of its graded Karoubi envelope, viewed as a  $\mathbb{Z}[q, q^{-1}]$ -algebra with the action of  $q$  coming from the grading shift functor, is isomorphic to  ${}_{\mathbb{Z}}\mathbf{U}_t^{\iota}$ , with the  $\iota$ -canonical basis arising from isomorphism classes of indecomposable objects.

The main theorem about  $\mathcal{N}\mathcal{B}_t$  proved in the present article gives explicit bases for morphism spaces in  $\mathcal{N}\mathcal{B}_t$ , a result which is used in an essential way in [11]. Our proof follows a similar strategy to the approach developed for Khovanov's Heisenberg category in [10], exploiting a remarkable monoidal functor

$$\Omega_t : \mathcal{N}\mathcal{B}_t \longrightarrow \text{Add}\left(\mathcal{U}(\mathfrak{sl}_2; t)_{\circ\dots\circ}\right)$$

from  $\mathcal{N}\mathcal{B}_t$  to the additive envelope of a monoidal category  $\mathcal{U}(\mathfrak{sl}_2; t)_{\circ\dots\circ}$  obtained by localizing the 2-category  $\mathcal{U}(\mathfrak{sl}_2)$  at certain morphisms. This functor takes the generating object  $B$  of  $\mathcal{N}\mathcal{B}_t$  to the direct sum  $E \oplus F$  of the generating objects of  $\mathcal{U}(\mathfrak{sl}_2; t)_{\circ\dots\circ}$ . It can be interpreted (in a weak sense due to the need to localize) as a categorification of the natural embedding of  ${}_{\mathbb{Z}}\mathbf{U}_t^{\iota}$  into the completion  $\prod_{\lambda, \mu \equiv t \pmod{2}} 1_{\mu} {}_{\mathbb{Z}}\mathbf{U} 1_{\lambda}$  of Lusztig's modified  $\mathbb{Z}[q, q^{-1}]$ -form of the quantum group  $U_q(\mathfrak{sl}_2)$ . Although the construction of  $\Omega_t$  is elementary, it depends on an astonishingly delicate computation with generators and relations expressed in terms of string calculus.

The rest of the article is organized as follows.

- We begin in Section 2 by defining  $\mathcal{N}\mathcal{B}_t$  by generators and relations; see Definition 2.1. Then we show that it is trivial unless  $t \in \{0, 1\}$ . For these values of  $t$ , we construct a homomorphism

$$\gamma_t : \Gamma \rightarrow \text{End}_{\mathcal{N}\mathcal{B}_t}(\mathbb{1})$$

where  $\Gamma$  is the subalgebra of the algebra of symmetric functions over  $\mathbb{k}$  generated by Schur's  $Q$ -functions.

- In Section 3, we recall the definition of the 2-category  $\mathcal{U}(\mathfrak{sl}_2)$ , and then introduce a localized version of it, denoted  $\mathcal{U}(\mathfrak{sl}_2)_{\circ\dots\circ}$ . In particular, this localization adjoins inverses of the 2-morphisms usually denoted by dots. We then derive some remarkable relations in  $\mathcal{U}(\mathfrak{sl}_2)_{\circ\dots\circ}$  involving certain 2-morphisms which we call *internal bubbles* which are analogous to, but more complicated than, corresponding morphisms for the Heisenberg category introduced in [10].
- Then in Section 4 we pass from the 2-category  $\mathcal{U}(\mathfrak{sl}_2)_{\circ\dots\circ}$  to the monoidal category  $\mathcal{U}(\mathfrak{sl}_2; t)_{\circ\dots\circ}$ . The objects in the latter are words  $X$  in  $E$  and  $F$ , corresponding to formal sums of the 1-morphisms  $X 1_{\lambda}$  in  $\mathcal{U}(\mathfrak{sl}_2)_{\circ\dots\circ}$  for the same word  $X$  and all weights  $\lambda \equiv t \pmod{2}$ , and its morphisms are represented by sequences of 2-morphisms in  $\mathcal{U}(\mathfrak{sl}_2)_{\circ\dots\circ}$ ; see Definition 4.1 for the complete definition. The monoidal functor  $\Omega_t$  is finally constructed in Theorem 4.2.
- In Section 5, we use this functor to prove our main basis theorem, Theorem 5.1. This result is analogous to the basis theorem establishing non-degeneracy of the Kac-Moody 2-category  $\mathcal{U}(\mathfrak{sl}_n)$  in [15]. It shows that the homomorphism  $\gamma_t$  mentioned earlier is actually an *isomorphism*, i.e.,  $\text{End}_{\mathcal{N}\mathcal{B}_t}(\mathbb{1}) \cong \Gamma$ . Moreover, each morphism space  $\text{Hom}_{\mathcal{N}\mathcal{B}_t}(B^{*n}, B^{*m})$  is free as a graded  $\Gamma$ -module with an explicit homogeneous basis.

Now we can give some further justification of the importance of the basis theorem proved here: in the sequel [11], we show that the graded rank of  $\text{Hom}_{\mathcal{NB}_t}(B^{*n}, B^{*m})$  as a free  $\Gamma$ -module is equal to  $(B^n, B^m)^t$ , where  $(\cdot, \cdot)^t$  is the non-degenerate symmetric bilinear form on the  $t$ -quantum group  ${}_{\mathbb{Z}}\mathbf{U}_t^i$  from [2, Lemma 6.25]. We regard this as the first indication that  $\mathcal{NB}_t$  categorifies  ${}_{\mathbb{Z}}\mathbf{U}_t^i$ , indeed, it provided us with some initial clues as to the precise form of the generators and relations for  $\mathcal{NB}_t$  in Definition 2.1.

2. DEFINITION AND FIRST PROPERTIES OF THE NIL-BRAUER CATEGORY

The definition of  $\mathcal{NB}_t$  is by generators and relations. We use the string calculus to denote morphisms in strict monoidal categories (or 2-morphisms in strict 2-categories), our general convention being that composition  $f \circ g$  (“vertical composition”) is depicted by placing  $f$  on top of  $g$  and tensor product  $f \star g$  (“horizontal composition”) is depicted by placing  $f$  to the left of  $g$ .

**Definition 2.1.** For  $t \in \mathbb{k}$ , the *nil-Brauer category*  $\mathcal{NB}_t$  is the strict monoidal category with one generating object  $B$ , whose identity endomorphism will be represented diagrammatically by the unlabeled string  $|$ , and four generating morphisms

$$\begin{array}{cccc}
 \bullet : B \rightarrow B, & \times : B \star B \rightarrow B \star B, & \cap : B \star B \rightarrow 1, & \cup : 1 \rightarrow B \star B, \\
 \text{(degree 2)} & \text{(degree -2)} & \text{(degree 0)} & \text{(degree 0)}
 \end{array} \tag{2.1}$$

subject to the following relations:

$$\begin{array}{c} \text{crossing} \end{array} = 0, \qquad \begin{array}{c} \text{crossing} \end{array} = \begin{array}{c} \text{crossing} \end{array}, \tag{2.2}$$

$$\text{circle} = t1_{\mathbb{1}}, \qquad \begin{array}{c} \cup \\ \cap \end{array} = | = \begin{array}{c} \cup \\ \cap \end{array}, \tag{2.3}$$

$$\begin{array}{c} \text{loop} \end{array} = 0, \qquad \begin{array}{c} \cap \\ \cup \end{array} = \begin{array}{c} \cup \\ \cap \end{array}, \tag{2.4}$$

$$\begin{array}{c} \bullet \\ \times \end{array} - \begin{array}{c} \times \\ \bullet \end{array} = | - \begin{array}{c} \cup \\ \cap \end{array}, \qquad \begin{array}{c} \cap \\ \bullet \end{array} = - \begin{array}{c} \bullet \\ \cap \end{array}. \tag{2.5}$$

**Remark 2.2.** Although it will only play a secondary role in this article, it is important for the sequel to note that  $\mathcal{NB}_t$  can be viewed as a *graded* monoidal category, i.e., a monoidal category enriched in the closed symmetric monoidal category of graded  $\mathbb{k}$ -modules, by declaring that the generators are of the degrees indicated in parentheses in (2.1).

**Remark 2.3.** The defining relations just recorded are quite familiar in related settings. The first two relations (2.2) are the same as defining relations in the nil-Hecke algebras associated to symmetric groups, but the polynomial generator of the nil-Hecke algebra (often depicted also by a dot) satisfies slightly different relations to (2.5). These relations come instead from the defining relations for the *affine Brauer category* of [21], which is the monoidal category defined in the same way as in Definition 2.1 but replacing the 0 on the right-hand side of the first relation in (2.2) by the identity (so that the crossing is an involution) and the 0 on the right-hand side of the first relation in (2.4) by  $\cap$ . The endomorphism algebras of objects in the affine Brauer category were introduced earlier by Nazarov [19, Section 4], and are known as Nazarov–Wenzl algebras [1], degenerate affine BMW algebras [13] or affine VW algebras [14]. The critical sign in the final relation

from (2.5) emerged in that setting from considerations involving orthogonal groups (and is unrelated to superalgebra).

As usual with definitions by generators and relations, the first task is to derive further relations as consequences of the defining ones. To start with, we have the following:

$$\begin{array}{c} \text{cup} = \text{uncup}, \end{array} \quad \begin{array}{c} \text{bubble} = 0 = \text{cap}, \end{array} \quad (2.6)$$

$$\begin{array}{c} \text{figure-eight} = 0, \end{array} \quad \begin{array}{c} \text{crossing} = 0, \end{array} \quad (2.7)$$

$$\begin{array}{c} \text{dot-crossing} - \text{dot-crossing} = \text{dot-line} - \text{dot-line}, \end{array} \quad \begin{array}{c} \text{dot-cup} = - \text{dot-cap}. \end{array} \quad (2.8)$$

For example, to prove the first of these, we attach cups to the bottom left and bottom right of the second relation in (2.4) to obtain

$$\text{cup} \cup \text{cup} = \text{cup} \cup \text{cup}.$$

This can then be simplified using the zigzag relations from (2.3) to obtain the desired relation. The other relations in (2.6) to (2.8) are proved similarly by attaching cups and/or caps to the ends of the strings in other defining relations then simplifying in obvious ways.

Now we take the first relation from (2.5) and close at the top by attaching a cap and at the bottom by attaching a cup. The left-hand side is 0 due to the first relations from (2.4) and (2.7). After replacing the bubbles on the right-hand side with  $t1_{\mathbb{1}}$ , we deduce from this that

$$t^2 1_{\mathbb{1}} = t 1_{\mathbb{1}}. \quad (2.9)$$

Thus, for  $\mathcal{NB}_t$  to be non-trivial, one must have that  $t \in \{0, 1\}$ . This will be assumed from now on. In fact, henceforth, we will treat  $t$  as though it is an element of  $\{0, 1\} \subset \mathbb{Z}$  (rather than being the image of that integer in  $\mathbb{k}$ ) so that we can use convenient expressions like  $(-1)^t$ .

The relations established so far imply that there are strict monoidal functors

$$R : \mathcal{NB}_t \rightarrow \mathcal{NB}_t^{\text{rev}}, \quad B \mapsto B, \quad s \mapsto (-1)^{\bullet(s)} s^{\leftrightarrow}, \quad (2.10)$$

$$T : \mathcal{NB}_t \rightarrow \mathcal{NB}_t^{\text{op}}, \quad B \mapsto B, \quad s \mapsto s^{\updownarrow}. \quad (2.11)$$

Here, for a string diagram  $s$  we use  $s^{\updownarrow}$  and  $s^{\leftrightarrow}$  to denote its reflection in a horizontal or vertical axis, and  $\bullet(s)$  denotes the total number of dots in the diagram. The category  $\mathcal{NB}_t$  is *strictly pivotal* with duality functor  $D := R \circ T = T \circ R$ ; this rotates a string diagram  $s$  through  $180^\circ$  then scales by  $(-1)^{\bullet(s)}$ . Also by the relations established so far, a string diagram with no dots can be deformed freely under planar isotopy without changing the morphism that it represents. This is not true in the presence of dots due to the sign in the last relations from (2.5) and (2.8)—there is a sign change whenever a dot slides across the critical point of a cup or cap.

To establish additional useful relations, we adopt a generating function formalism which is a slight refinement of the setup introduced in [8]. We denote the  $r$ th power of  $\bullet$  under vertical composition simply by labeling the dot with  $r$ . More generally, given a polynomial  $f(x) = \sum_{r \geq 0} c_r x^r \in \mathbb{k}[x]$  and a dot in some string diagram  $s$ , we denote

$$\sum_{r \geq 0} c_r \times (\text{the morphism obtained from } s \text{ by labeling the dot by } r)$$

by attaching what we call a *pin* to the dot, labeling the node at the head of the pin by  $f(x)$ :

$$\bullet \text{---} \boxed{f(x)} := \sum_{r \geq 0} c_r \bullet \text{---} r \in \text{End}_{\mathcal{NB}_t}(B). \tag{2.12}$$

In the drawing of a pin, the arm and the head of the pin can be moved freely around larger diagrams so long as the point stays put—these are not part of the string calculus. More generally,  $f(x)$  here could be a polynomial with coefficients in the algebra  $\mathbb{k}((u^{-1}))$  of formal Laurent series in an indeterminate  $u^{-1}$ ; then the string  $s$  decorated with a pin labeled  $f(x)$  defines a morphism in the base-changed monoidal category  $\mathcal{NB}_t[[u^{-1}]]$ . We think of this as being a generating function for a family of morphisms. Pins labeled by the power series

$$(u + ax)^{-1} = u^{-1} - axu^{-2} + a^2x^2u^{-3} - a^3x^3u^{-4} + \dots \in \mathbb{k}[x][[u^{-1}]] \tag{2.13}$$

for  $a \in \{+, -\}$  appear so often that we denote them by a special shorthand, putting the variable  $u$  into the node at the head of the pin and in addition we label the arm of the pin by the sign  $a$ :

$$\begin{aligned} & \bullet \text{---} \overset{-}{\circlearrowleft} u \\ & := \bullet \text{---} \boxed{(u-x)^{-1}} = u^{-1} \left| + u^{-2} \bullet \text{---} + u^{-3} \bullet \text{---} \bullet \text{---} + u^{-4} \bullet \text{---} \bullet \text{---} \bullet \text{---} + \dots \in \text{End}_{\mathcal{NB}_t}(B)[[u^{-1}]], \end{aligned} \tag{2.14}$$

$$\begin{aligned} & \bullet \text{---} \overset{+}{\circlearrowright} u \\ & := \bullet \text{---} \boxed{(u+x)^{-1}} = u^{-1} \left| - u^{-2} \bullet \text{---} + u^{-3} \bullet \text{---} \bullet \text{---} - u^{-4} \bullet \text{---} \bullet \text{---} \bullet \text{---} + \dots \in \text{End}_{\mathcal{NB}_t}(B)[[u^{-1}]]. \end{aligned} \tag{2.15}$$

These shorthand symbols behave well under  $R$  and  $T$  (2.10) and (2.11):

$$R \left( \bullet \text{---} \overset{a}{\circlearrowleft} u \right) = \overset{-a}{\circlearrowright} u \bullet, \quad T \left( \bullet \text{---} \overset{a}{\circlearrowright} u \right) = \bullet \text{---} \overset{a}{\circlearrowleft} u. \tag{2.16}$$

**Lemma 2.4.** *The following relations hold in  $\mathcal{NB}_t$ :*

$$\left( \bullet \text{---} \overset{a}{\circlearrowleft} u \right) = \left( \overset{-a}{\circlearrowright} u \bullet \right), \tag{2.17}$$

$$\left( \bullet \text{---} \overset{a}{\circlearrowright} u \right) = \left( \overset{-a}{\circlearrowleft} u \bullet \right), \tag{2.18}$$

$$\left( \begin{array}{c} \diagdown \bullet \text{---} \overset{a}{\circlearrowleft} u \diagup \\ \bullet \text{---} \overset{a}{\circlearrowleft} u \end{array} \right) - \left( \begin{array}{c} \bullet \text{---} \overset{a}{\circlearrowleft} u \\ \diagdown \bullet \text{---} \overset{a}{\circlearrowleft} u \diagup \end{array} \right) = a \left( \begin{array}{c} \bullet \text{---} \overset{a}{\circlearrowleft} u \\ \bullet \text{---} \overset{a}{\circlearrowleft} u \end{array} \right) - a \left( \begin{array}{c} \bullet \text{---} \overset{a}{\circlearrowleft} u \\ \bullet \text{---} \overset{a}{\circlearrowleft} u \end{array} \right), \tag{2.19}$$

$$\left( \begin{array}{c} \bullet \text{---} \overset{a}{\circlearrowright} u \\ \diagdown \bullet \text{---} \overset{a}{\circlearrowright} u \diagup \end{array} \right) - \left( \begin{array}{c} \diagdown \bullet \text{---} \overset{a}{\circlearrowright} u \diagup \\ \bullet \text{---} \overset{a}{\circlearrowright} u \end{array} \right) = a \left( \begin{array}{c} \bullet \text{---} \overset{a}{\circlearrowright} u \\ \bullet \text{---} \overset{a}{\circlearrowright} u \end{array} \right) - a \left( \begin{array}{c} \bullet \text{---} \overset{a}{\circlearrowright} u \\ \bullet \text{---} \overset{a}{\circlearrowright} u \end{array} \right). \tag{2.20}$$

*Proof.* It is clear from the last relation in (2.5) that  $\left( \bullet \text{---} \boxed{f(x)} \right) = \left( \boxed{f(-x)} \bullet \right)$  and similarly for cups. The relations (2.17) and (2.18) follow. To prove (2.19), it suffices to establish the equivalent relation obtained by vertically composing on top with  $\bullet \text{---} \overset{+}{\circlearrowright} u$  and on the bottom with  $\bullet \text{---} \overset{-}{\circlearrowleft} u$ ; this equivalent relation follows immediately from the first relation from (2.5). Finally, the relation (2.20) follows on applying  $T$ .  $\square$

The pin notation gives the generating function

$$\bullet \text{---} \overset{-}{\circlearrowleft} u = \sum_{r \geq 0} \bullet \text{---} r u^{-r-1} \in \text{End}_{\mathcal{NB}_t}(\mathbb{1})[[u^{-1}]] \tag{2.21}$$

for “dotted bubbles”. This is often useful, but even more important is the renormalization

$$\mathbb{O}(u) = \sum_{r \geq 0} \mathbb{O}_r u^{-r} := (-1)^t \left( 1_{\mathbb{1}} - 2u \text{ (dotted bubble)} \right) \in 1_{\mathbb{1}} + u^{-1} \text{End}_{\mathcal{NB}_t}(\mathbb{1})[[u^{-1}]]. \quad (2.22)$$

Its coefficients are given explicitly by

$$\mathbb{O}_0 := 1_{\mathbb{1}}, \quad \mathbb{O}_r := -2(-1)^t \text{ (dotted bubble)}^r \quad (2.23)$$

for  $r \geq 1$ . Note also by (2.16) and (2.17) that  $\mathbb{O}(u)$  is invariant under both of the symmetries  $\mathbb{R}$  and  $\mathbb{T}$ .

The following derives some further relations involving these generating functions. Yet more can then be obtained by applying  $\mathbb{R}$  and  $\mathbb{T}$ .

**Theorem 2.5.** *The following relations hold in  $\mathcal{NB}_t$ :*

$$2u \text{ (dotted bubble)} = 2u \left( \text{ (dotted bubble)}^+ - \text{ (dotted bubble)}^- \right), \quad (2.24)$$

$$\text{ (dotted bubble)}^- + \text{ (dotted bubble)}^+ = 2u \text{ (dotted bubble)}^+ \text{ (dotted bubble)}^-, \quad (2.25)$$

$$\mathbb{O}(u)\mathbb{O}(-u) = 1_{\mathbb{1}}, \quad (2.26)$$

$$\mathbb{O}(u) \Big| = \left( \frac{u-x}{u+x} \right)^2 \text{ (dotted bubble)} \mathbb{O}(u). \quad (2.27)$$

*Proof.* To prove (2.24), we have that

$$\begin{aligned} 2u \text{ (dotted bubble)} &\stackrel{(2.18)}{=} 2u \text{ (dotted bubble)}^+ \\ &\stackrel{(2.19)}{=} 2u \left( \text{ (dotted bubble)}^+ - \text{ (dotted bubble)}^- \right) \\ &= 2u \left( \text{ (dotted bubble)}^+ \text{ (dotted bubble)}^- - \text{ (dotted bubble)}^- \text{ (dotted bubble)}^+ \right) \\ &= 2u \left( \text{ (dotted bubble)}^+ \text{ (dotted bubble)}^- - \text{ (dotted bubble)}^- \text{ (dotted bubble)}^+ \right). \end{aligned}$$

The desired relation (2.24) follows easily from this using  $\frac{2u}{(u+x)(u-x)} = \frac{1}{u-x} + \frac{1}{u+x}$ .

Closing the free strings on the left in (2.24) gives a new relation whose left-hand side is zero. This gives (2.25) immediately. Then (2.26) follows by substituting the definition of  $\mathbb{O}(u)$  and  $\mathbb{O}(-u)$  into the product  $\mathbb{O}(u)\mathbb{O}(-u)$ , multiplying out the brackets, and simplifying using (2.25).

Finally, for the “bubble slide” relation (2.27), we have that

$$0 = \text{ (dotted bubble)}^+ \text{ (dotted bubble)}^- = \text{ (dotted bubble)}^- \text{ (dotted bubble)}^+ = \text{ (dotted bubble)}^+ \text{ (dotted bubble)}^- - \text{ (dotted bubble)}^- \text{ (dotted bubble)}^+.$$

This shows that

$$\text{ (dotted bubble)}^+ \text{ (dotted bubble)}^- = \text{ (dotted bubble)}^- \text{ (dotted bubble)}^+.$$

Let  $f$  denote the morphism on the left-hand side of this equation. The equation shows that  $f$  is fixed by the symmetry  $\mathbb{R}$ . Expanding the curl using (2.24) gives that

$$2uf = 2u \left( \text{ (dotted bubble)}^+ \text{ (dotted bubble)}^- - \text{ (dotted bubble)}^- \text{ (dotted bubble)}^+ \right).$$



So, using the definition of  $\mathbb{O}(u)$ , we have that

$$-(-1)^t \left( 2uf + \begin{array}{c} \bullet \\ \mid \\ \ominus u \\ \mid \\ \bullet \\ \oplus u \\ \mid \\ \bullet \end{array} \right) = \begin{array}{c} \bullet \\ \mid \\ \oplus u \\ \mid \\ \bullet \\ \oplus u \\ \mid \\ \bullet \end{array} \times (-1)^t \left( 1_{\mathbb{1}} - 2u \begin{array}{c} \bullet \\ \mid \\ \ominus u \\ \mid \\ \bullet \end{array} \right) = \boxed{(u+x)^{-2}} \begin{array}{c} \bullet \\ \mid \\ \bullet \end{array} \mathbb{O}(u).$$

Since  $f = \mathbb{R}(f)$ , the expression on the left-hand side of this equation is fixed by  $\mathbb{R}$ , hence, so is the expression on the right-hand side. Since  $\mathbb{R}(\mathbb{O}(u)) = \mathbb{O}(u)$ , this implies that

$$\boxed{(u+x)^{-2}} \begin{array}{c} \bullet \\ \mid \\ \bullet \end{array} \mathbb{O}(u) = \mathbb{O}(u) \begin{array}{c} \bullet \\ \mid \\ \bullet \end{array} \boxed{(u-x)^{-2}}.$$

The relation (2.27) follows immediately from this. □

Let  $\Lambda$  be the algebra of symmetric functions over the ground ring  $\mathbb{k}$ . Adopting standard notations, this is freely generated either by the elementary symmetric functions  $e_r$  ( $r > 0$ ) or by the complete symmetric functions  $h_r$  ( $r > 0$ ). The two families of generators are related by the identity

$$e(-u)h(u) = 1 \tag{2.28}$$

where

$$e(u) = \sum_{r \geq 0} e_r u^{-r}, \quad h(u) = \sum_{r \geq 0} h_r u^{-r} \tag{2.29}$$

are the corresponding generating functions, and  $e_0 = h_0 = 1$  by convention.

Following [18, Chapter III, Section 8], we define  $q(u) \in \Lambda[[u^{-1}]]$  and elements  $q_r$  ( $r \geq 0$ ) of  $\Lambda$  so that

$$q(u) = \sum_{r \geq 0} q_r u^{-r} := e(u)h(u). \tag{2.30}$$

By (2.28), we have that

$$q(u)q(-u) = 1 \tag{2.31}$$

Equivalently,  $q_0 = 1$  and

$$q_{2r} = (-1)^{r-1} \frac{1}{2} q_r^2 + \sum_{s=1}^{r-1} (-1)^{s-1} q_s q_{2r-s} \tag{2.32}$$

for  $r \geq 1$ ; cf. [18, (III.8.2')]. The subalgebra of  $\Lambda$  generated by all  $q_r$  ( $r \geq 0$ ) is denoted  $\Gamma$ . As explained in [18],  $\Gamma$  is freely generated by  $q_1, q_3, q_5, \dots$  (and it has a distinguished basis given by the Schur  $Q$ -functions  $Q_\lambda$  indexed by all strict partitions). It follows that  $\Gamma$  is generated by the elements  $q_r$  ( $r \geq 0$ ) subject just to the relations (2.31). Hence, the relation (2.26) from Theorem 2.5 implies the following:

**Corollary 2.6.** *There is a unique algebra homomorphism  $\gamma_t : \Gamma \rightarrow \text{End}_{\mathcal{N}\mathbb{B}_t}(\mathbb{1})$  such that  $q_r \mapsto \mathbb{O}_r$  for all  $r \geq 0$ .*

We will show in Corollary 5.4 below that the homomorphism  $\gamma_t$  just constructed is actually an *isomorphism*.

### 3. RELATIONS IN THE 2-CATEGORY $\mathfrak{U}(\mathfrak{sl}_2)$

Next we recall the definition of the 2-category  $\mathfrak{U}(\mathfrak{sl}_2)$  following the approach of Lauda [16]. Working still over the ground ring  $\mathbb{k}$ ,  $\mathfrak{U}(\mathfrak{sl}_2)$  is the strict 2-category with object set  $\mathbb{Z}$ , generating 1-morphisms  $E1_\lambda : \lambda \rightarrow \lambda + 2$  and  $1_\lambda F : \lambda + 2 \rightarrow \lambda$ , whose identity 2-morphisms will be represented by the oriented strings  $\uparrow_\lambda$  and  $\downarrow_\lambda$ , and generating 2-morphisms

$$\begin{aligned} \uparrow_\lambda : E1_\lambda \Rightarrow E1_\lambda, & \quad \curvearrowright_\lambda : EF1_\lambda \Rightarrow 1_\lambda, & \quad \cup_\lambda : 1_\lambda \Rightarrow FE1_\lambda, & \quad (3.1) \\ \text{(degree 2)} & \quad \text{(degree } 1 - \lambda) & \quad \text{(degree } \lambda + 1) \end{aligned}$$

$$\begin{aligned} \times_\lambda : EE1_\lambda \Rightarrow EE1_\lambda, & \quad \curvearrowleft_\lambda : FE1_\lambda \Rightarrow 1_\lambda, & \quad \cup_\lambda : 1_\lambda \Rightarrow EF1_\lambda, & \quad (3.2) \\ \text{(degree } -2) & \quad \text{(degree } \lambda + 1) & \quad \text{(degree } 1 - \lambda) \end{aligned}$$

for all  $\lambda \in \mathbb{Z}$ , subject to certain relations below. We refer to the dots in this setting as *open dots* to distinguish them from the (closed) dots in the nil-Brauer category in the previous section.

**Remark 3.1.** The 2-category  $\mathfrak{U}(\mathfrak{sl}_2)$  also admits a grading making it into a strict graded 2-category. This is defined by declaring that the generating 2-morphisms are of the degrees specified in parentheses. This grading will usually be ignored, but it crops up again in Remark 4.4, and it will be needed in the final section since we need there to pass to the completion of  $\mathfrak{U}(\mathfrak{sl}_2)$  with respect to the grading.

To write down the relations, we denote the  $r$ th power of the dot under vertical composition simply by labeling it with the natural number  $r$ . We introduce rightward and leftward crossings by setting

$$\begin{aligned} \times_\lambda &:= \text{rightward crossing}, & \times_\lambda &:= \text{leftward crossing}, & \quad (3.3) \\ \text{(degree 0)} & & \text{(degree 0)} & \end{aligned}$$

and use the following shorthands:

$$\begin{aligned} \circlearrowleft_n \lambda &:= \begin{cases} (-1)^n \det \left( \lambda \circlearrowleft_{r-s-\lambda} \right)_{r,s=1,\dots,n} & \text{if } 0 \leq n \leq -\lambda \\ 0 & \text{if } n < 0 \text{ or } n > -\lambda, \end{cases} & \quad (3.4) \\ \text{(degree } 2n) & \end{aligned}$$

$$\begin{aligned} \lambda \circlearrowright_n &:= \begin{cases} (-1)^n \det \left( \lambda \circlearrowright_{r-s+\lambda} \right)_{r,s=1,\dots,n} & \text{if } 0 \leq n \leq \lambda \\ 0 & \text{if } n < 0 \text{ or } n > \lambda. \end{cases} & \quad (3.5) \\ \text{(degree } 2n) & \end{aligned}$$

Then the defining relations are as follows:

$$\begin{aligned} \text{cup} \cap &= 0, & \text{rightward crossing} &= \text{leftward crossing}, & \text{rightward crossing} - \text{leftward crossing} &= \text{rightward crossing} - \text{leftward crossing} = \uparrow_\lambda - \downarrow_\lambda, & \quad (3.6) \end{aligned}$$

$$\begin{aligned} \text{cup} &= \uparrow_\lambda, & \text{cap} &= \downarrow_\lambda, & \quad (3.7) \end{aligned}$$

$$\begin{aligned}
 \begin{array}{c} \text{crossing} \\ \lambda \end{array} &= - \begin{array}{c} \text{vertical line} \\ \lambda \end{array} + \sum_{n=0}^{-\lambda-1} \sum_{\substack{r,s \geq 0 \\ r+s = -\lambda-1-n}} \begin{array}{c} \text{cup with dots} \\ \lambda \end{array}, \\
 \begin{array}{c} \text{crossing} \\ \lambda \end{array} &= - \begin{array}{c} \text{vertical line} \\ \lambda \end{array} + \sum_{n=0}^{\lambda-1} \sum_{\substack{r,s \geq 0 \\ r+s = \lambda-1-n}} \begin{array}{c} \text{cup with dots} \\ \lambda \end{array},
 \end{aligned} \tag{3.8}$$

$$\begin{array}{c} \text{loop} \\ \lambda \end{array} = -\delta_{\lambda,0} \begin{array}{c} \text{vertical line} \\ \lambda \end{array} \text{ if } \lambda \geq 0, \quad \begin{array}{c} \text{loop} \\ \lambda \end{array} = \delta_{\lambda,0} \begin{array}{c} \text{vertical line} \\ \lambda \end{array} \text{ if } \lambda \leq 0, \tag{3.9}$$

$$\begin{array}{c} \text{loop with dot} \\ \lambda \end{array} = \delta_{n,\lambda-1} 1_{1_\lambda} \text{ for all } 0 \leq n < \lambda, \quad \begin{array}{c} \text{loop with dot} \\ \lambda \end{array} = \delta_{n,-\lambda-1} 1_{1_\lambda} \text{ for all } 0 \leq n < -\lambda. \tag{3.10}$$

A more efficient presentation for  $\mathfrak{U}(\mathfrak{sl}_2)$  was given by Rouquier in [20]. To formulate this, we just need the generating morphisms  $\hat{\phi}_\lambda, \hat{\chi}_\lambda, \cup_\lambda$  and  $\cap_\lambda$  (hence, we also have the rightward crossings defined via (3.3)), subject to the relations (3.6), the zigzag identities from (3.7), and the new *inversion relation* which asserts that the following matrices are invertible in the additive envelope of  $\mathfrak{U}(\mathfrak{sl}_2)$ :

$$\left[ \hat{\chi}_\lambda, \cup_{\lambda-1}, \dots, \cup_\lambda \right] : EF1_\lambda \oplus 1_\lambda^{\oplus -\lambda} \Rightarrow FE1_\lambda \quad \text{if } \lambda \leq 0, \tag{3.11}$$

$$\begin{bmatrix} \hat{\chi}_\lambda \\ \cap_{\lambda-1} \\ \vdots \\ \cap_\lambda \end{bmatrix} : EF1_\lambda \Rightarrow FE1_\lambda \oplus 1_\lambda^{\oplus \lambda} \quad \text{if } \lambda \geq 0. \tag{3.12}$$

The equivalence of Rouquier’s presentation with Lauda’s one from the previous paragraph was established in [5]. It is shown there that the two-sided inverse of the matrix (3.11) is the  $(1 - \lambda) \times 1$  matrix with first entry  $-\hat{\chi}_\lambda$  and  $(n + 2)^{\text{th}}$  entry equal to  $\lambda \cap_{\circ n} + \lambda \cap_{\circ n-1} \textcircled{1} + \dots + \lambda \cap_{\circ n}$  for  $0 \leq n \leq -\lambda - 1$ , and the two-sided inverse of the matrix (3.12) is the  $1 \times (\lambda + 1)$  matrix with first entry  $-\hat{\chi}_\lambda$  and  $(n + 2)^{\text{th}}$  entry equal to  $n \hat{\phi}_\cup \lambda + \textcircled{1} n-1 \hat{\phi}_\cup \lambda + \dots + \textcircled{n} \cup_\lambda$  for  $0 \leq n \leq \lambda - 1$ .

We define the downward open dot  $\lambda \downarrow$  and the downward crossing  $\lambda \searrow$  to be the right mates of  $\hat{\phi}_\lambda$  and  $\hat{\chi}_\lambda$ , respectively. The defining relations for  $\mathfrak{U}(\mathfrak{sl}_2)$  imply that

$$\begin{array}{c} \text{cup} \\ \lambda \end{array} = \begin{array}{c} \text{vertical line} \\ \lambda \end{array}, \quad \begin{array}{c} \text{cup} \\ \lambda \end{array} = \begin{array}{c} \text{vertical line} \\ \lambda \end{array}. \tag{3.13}$$

Moreover,  $\lambda \downarrow$  and  $\lambda \searrow$  are equal to the *left* mates of  $\hat{\phi}_\lambda$  and  $\hat{\chi}_\lambda$ . It follows that diagrams for 2-morphisms in  $\mathfrak{U}(\mathfrak{sl}_2)$  are invariant under boundary-preserving planar isotopy. In particular,  $\mathfrak{U}(\mathfrak{sl}_2)$  is strictly pivotal with duality functor  $\mathbb{D}$  defined by rotating diagrams through  $180^\circ$ . There are a couple more useful symmetries, i.e., strict 2-functors

$$\mathbb{R} : \mathfrak{U}(\mathfrak{sl}_2) \rightarrow \mathfrak{U}(\mathfrak{sl}_2)^{\text{rev}}, \quad \mathbb{T} : \mathfrak{U}(\mathfrak{sl}_2) \rightarrow \mathfrak{U}(\mathfrak{sl}_2)^{\text{op}} \tag{3.14}$$

defined as follows:

- $R$  takes the object  $\lambda$  to  $-\lambda$ , switches  $E1_\lambda$  with  $1_{-\lambda}E$  and  $F1_\lambda$  with  $1_{-\lambda}F$ , and takes string diagram  $s$  to  $(-1)^{\times(s)}s^{\leftrightarrow}$ , also negating weights  $\lambda$  labeling regions;
- $T$  takes  $\lambda$  to  $-\lambda$ , switches the generating 1-morphisms  $E1_\lambda$  and  $F1_{-\lambda}$ , and takes string diagram  $s$  representing a 2-morphism to  $(-1)^{\times(s)}s^\updownarrow$ , also negating weights  $\lambda$  labeling regions.

Here,  $\times(s)$  is the total number of crossings in  $s$ . The duality functor factorizes as  $D = R \circ T = T \circ R$ .

We use similar conventions for pins attached to open dots as in (2.12) for pins attached to closed dots in the previous section. In particular, as in (2.14) and (2.15), we have

$$\begin{aligned} \uparrow \text{---} \textcircled{-u} \lambda &:= \uparrow \text{---} \boxed{(u-x)^{-1}} \lambda = u^{-1} \uparrow \lambda + u^{-2} \uparrow \textcircled{\lambda} + u^{-3} \uparrow \textcircled{\textcircled{\lambda}} + \dots, & (3.15) \\ \uparrow \text{---} \textcircled{+u} \lambda &:= \uparrow \text{---} \boxed{(u+x)^{-1}} \lambda = u^{-1} \uparrow \lambda - u^{-2} \uparrow \textcircled{\lambda} + u^{-3} \uparrow \textcircled{\textcircled{\lambda}} - \dots. & (3.16) \end{aligned}$$

Now, unlike in (2.17) and (2.18), we have simply that

$$\begin{aligned} \textcircled{\textcircled{a}} \text{---} \textcircled{u} &= \textcircled{u} \text{---} \textcircled{a}, & \textcircled{a} \text{---} \textcircled{u} &= \textcircled{u} \text{---} \textcircled{a}, & (3.17) \end{aligned}$$

and similarly for the other orientation.

We define the *fake bubble polynomials*

$$\begin{aligned} \textcircled{u} \lambda &:= \sum_{n=0}^{-\lambda} \textcircled{n} \lambda u^{-\lambda-n}, & \textcircled{u} \lambda &:= \sum_{n=0}^{\lambda} \textcircled{n} \lambda u^{\lambda-n}, & (3.18) \end{aligned}$$

which are polynomials in  $\text{End}_{\mathcal{U}(\mathfrak{sl}_2)}(1_\lambda)[u]$  with  $\textcircled{u} \lambda = \delta_{\lambda,0}1_{1_\lambda}$  when  $\lambda \geq 0$  and  $\textcircled{u} \lambda = \delta_{\lambda,0}1_{1_\lambda}$  when  $\lambda \leq 0$ . It is often convenient to combine the fake bubble polynomials with generating functions for genuinely dotted bubbles by letting

$$\lambda \textcircled{\circ}(u) := \textcircled{u} \lambda + \textcircled{u} \text{---} \textcircled{\circ} \lambda \in u^{-\lambda}1_{1_\lambda} + u^{-\lambda-1} \text{End}_{\mathcal{U}(\mathfrak{sl}_2)}(1_\lambda)[[u^{-1}]], \quad (3.19)$$

$$\lambda \textcircled{\circ}(u) := \textcircled{u} \lambda + \textcircled{\circ} \text{---} \textcircled{u} \lambda \in u^\lambda 1_{1_\lambda} + u^{\lambda-1} \text{End}_{\mathcal{U}(\mathfrak{sl}_2)}(1_\lambda)[[u^{-1}]]. \quad (3.20)$$

As explained originally in [17, Proposition 8.2] (see also [6, (3.11)–(3.12)]), for any  $\lambda \in \mathbb{Z}$ , the algebra  $\text{End}_{\mathcal{U}(\mathfrak{sl}_2)}(1_\lambda)$  may be identified with the algebra  $\Lambda$  of symmetric functions so that  $\lambda \textcircled{\circ}(u)$  and  $\lambda \textcircled{\circ}(u)$  as just defined are identified with the generating functions  $u^{-\lambda}e(-u)$  and  $u^\lambda h(u)$  from (2.29). In particular, we have that

$$\lambda \textcircled{\circ}(u) \textcircled{\circ}(u) = 1_{1_\lambda} \quad (3.21)$$

as in (2.28).

The following relations are proved in [5, Corollary 3.5]:

$$\begin{aligned} \textcircled{\circ} \lambda &= - \sum_{n=0}^{-\lambda} -\lambda-n \textcircled{n} \lambda = - \left[ \textcircled{-u} \textcircled{u} \lambda \right]_{u^{-1}}, & (3.22) \end{aligned}$$

$$\lambda \textcircled{\circ} = \sum_{n=0}^{\lambda} \lambda \textcircled{n} \textcircled{\circ} \lambda-n = \left[ \lambda \textcircled{u} \textcircled{-u} \right]_{u^{-1}}, \quad (3.23)$$

with the second equalities being easily checked by equating coefficients; here and below we use  $[f(u)]_{u^r}$  to denote the coefficient of  $u^r$  of a formal Laurent series  $f(u)$  in  $u^{-1}$ . Other

defining relations can be written similarly in terms of generating functions. For example, the following are equivalent to (3.8):

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \lambda = - \begin{array}{c} \uparrow \\ \downarrow \end{array} \lambda + \left[ \begin{array}{c} \text{bubble with } u \\ \text{bubble with } u \end{array} \right]_{u^{-1}}, \quad \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} \lambda = - \begin{array}{c} \uparrow \\ \downarrow \end{array} \lambda + \left[ \begin{array}{c} \text{bubble with } u \\ \text{bubble with } u \end{array} \right]_{u^{-1}}. \quad (3.24)$$

Next, we have the *bubble slide relations*

$$\begin{array}{c} \circlearrowleft(u) \uparrow \\ \downarrow \end{array} \lambda = \begin{array}{c} \text{bubble with } u \\ \text{bubble with } u \end{array} \circlearrowleft(u) \lambda, \quad \begin{array}{c} \uparrow \\ \circlearrowleft(u) \downarrow \end{array} \lambda = \begin{array}{c} \text{bubble with } u \\ \text{bubble with } u \end{array} \downarrow \lambda, \quad (3.25)$$

which follow from [15, Proposition 3.4]. They imply similar relations for the fake bubble polynomials:

$$\begin{array}{c} \text{bubble with } u \\ \uparrow \end{array} \lambda = \left[ \begin{array}{c} \text{bubble with } u \\ \text{bubble with } u \end{array} \begin{array}{c} \text{bubble with } u \\ \uparrow \end{array} \lambda \right]_{u \geq 0}, \quad \begin{array}{c} \uparrow \\ \text{bubble with } u \end{array} \lambda = \left[ \begin{array}{c} \text{bubble with } u \\ \text{bubble with } u \end{array} \begin{array}{c} \uparrow \\ \text{bubble with } u \end{array} \lambda \right]_{u \geq 0}. \quad (3.26)$$

Finally, we have the *alternating braid relation*

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \lambda - \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} \lambda = \left[ \begin{array}{c} \text{bubble with } u \\ \text{bubble with } u \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \lambda + \begin{array}{c} \downarrow \\ \uparrow \end{array} \begin{array}{c} \text{bubble with } u \\ \text{bubble with } u \end{array} \lambda \right]_{u^{-1}} \quad (3.27)$$

from [15, Proposition 3.5].

In order to make a connection between  $\mathfrak{U}(\mathfrak{sl}_2)$  and the nil-Brauer category, we need to localize at the morphisms

$$\begin{array}{c} \hat{\circ} \cdots \hat{\circ} \\ \text{---} \\ X \end{array} \hat{\circ} \lambda := \begin{array}{c} \hat{\circ} \\ \text{---} \\ X \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \lambda + \begin{array}{c} \uparrow \\ \text{---} \\ X \end{array} \hat{\circ} \lambda \quad (3.28)$$

for all  $\lambda, \mu \in \mathbb{Z}$  and all 1-morphisms  $X : \lambda + 2 \rightarrow \mu - 2$  in  $\mathfrak{U}(\mathfrak{sl}_2)$ . This means that we adjoin additional generating 2-morphisms  $\hat{\circ} \text{---} \hat{\circ} \lambda : EXE1_\lambda \Rightarrow EXE1_\lambda$  subject to the additional relations

$$\begin{array}{c} \hat{\circ} \text{---} \hat{\circ} \lambda \\ \text{---} \\ X \end{array} = \left( \begin{array}{c} \hat{\circ} \cdots \hat{\circ} \\ \text{---} \\ X \end{array} \hat{\circ} \lambda \right)^{-1}. \quad (3.29)$$

By some analogy with [9, (4.21)], we refer to the 2-morphisms (3.29) as *teleporters*<sup>1</sup>: the relation

$$\begin{array}{c} \hat{\circ} \text{---} \hat{\circ} \lambda \\ \text{---} \\ X \end{array} + \begin{array}{c} \hat{\circ} \text{---} \hat{\circ} \lambda \\ \text{---} \\ X \end{array} = \begin{array}{c} \uparrow \\ \text{---} \\ X \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \lambda \quad (3.30)$$

means that dots can “teleport” across teleporters (hence, the name!). We denote the strict 2-category obtained in this way by  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ \cdots \circ}$ ; it also admits a grading like in Remark 3.1.

In  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ \cdots \circ}$ , it is easy to see using (3.7) and (3.13) that the 2-morphisms defined similarly to (3.28) but with one or both of the upward strings changed to downward strings are also invertible; we denote their inverses in the obvious way by modifying the directions of arrows in (3.29). The dotted and solid horizontal lines in all of these diagrams

<sup>1</sup>They are also closely related to the morphisms called *dumbbells* in [10].

are present merely to indicate that the open dots at the endpoints have been identified—they are not a part of the string calculus so can be moved freely around larger diagrams as long as the endpoints remain fixed. It will often be convenient to allow open dots to be connected by dotted or solid lines even when the endpoints are not at the same horizontal level. Such string diagrams may be interpreted as morphisms by using planar isotopy to redraw the diagrams so that the endpoints are aligned. For example, we have that

$$\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \curvearrowright \lambda = \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \text{---} \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \cup \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda + \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \cup \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = 2 \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \cup \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = 2 \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda . \tag{3.31}$$

It is similarly straightforward to check that

$$\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \curvearrowleft \lambda = \left( \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \curvearrowright \lambda \right)^{-1} .$$

From (3.31), we deduce that open dots are invertible in  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ \dots \circ}$ . We will denote the inverse of  $\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda$  simply by labeling the dot by  $-1$ , and have that

$$\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \curvearrowleft \lambda = \frac{1}{2} \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \begin{array}{c} -1 \\ \uparrow \\ -1 \end{array} \lambda . \tag{3.32}$$

From (3.30), we deduce that

$$\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \text{---} \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda + \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \text{---} \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \begin{array}{c} -1 \\ \uparrow \\ -1 \end{array} \lambda . \tag{3.33}$$

We also have that

$$\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \text{---} \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda - \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \text{---} \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \begin{array}{c} -u \\ \uparrow \\ u \end{array} \lambda . \tag{3.34}$$

This and the following relations are easily deduced in a similar way to the proof of (2.19):

$$\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda - \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda - \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = -1 \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \begin{array}{c} -1 \\ \uparrow \\ -1 \end{array} \lambda , \tag{3.35}$$

$$\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda - \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda - \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = a \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \begin{array}{c} -a \\ \uparrow \\ a \end{array} \lambda , \tag{3.36}$$

$$\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda - \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda - \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \times \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda = \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda . \tag{3.37}$$

In the literature, the fake bubbles  $\begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda$  and  $\lambda \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array}$  are often denoted by  $-1 \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda$  and  $\lambda \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} -1$ , respectively. We have avoided the latter convention because in  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ \dots \circ}$  it makes sense to consider *genuine* dotted bubbles with dot labeled by  $-1$ .

**Lemma 3.2.**  $\left( -1 \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda - \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \lambda \right) \left( \lambda \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} -1 - \lambda \begin{array}{c} \circ \\ \uparrow \\ \circ \end{array} \right) = 1.$

*Proof.* By (3.35), we have that

$$\text{Diagram 1} - \text{Diagram 2} = \text{Diagram 3} - \text{Diagram 4} .$$

Then we expand the curls on the left-hand side using (3.22) and (3.23) to obtain

$$\sum_{n=0}^{\lambda} \text{Diagram 5} + \sum_{n=0}^{-\lambda} \text{Diagram 6} = \text{Diagram 7} .$$

By (3.10), the clockwise dotted bubble on the left-hand side here is 0 unless either  $n = 0 < \lambda$  (when it equals  $1_{1_\lambda}$ ) or  $n = \lambda$ , and the counterclockwise one is 0 unless either  $n = 0 < -\lambda$  (when it equals  $1_{1_\lambda}$ ) or  $n = -\lambda$ . Using this, and considering the cases  $\lambda > 0, \lambda = 0$  or  $\lambda < 0$  separately, the equation obtained so far reduces to the identity we are trying to prove.  $\square$

**Lemma 3.3.**  $\text{Diagram 8} - \text{Diagram 9} = \frac{1}{2} \text{Diagram 10} .$

*Proof.* Using the rotated version of the last relation from (3.6) to commute all the dots to the bottom, one checks easily that  $\text{Diagram 11} - \text{Diagram 12} = 2 \text{Diagram 13}$ . Now compose vertically with teleporters on the top and the bottom.  $\square$

We come to what we promise is the last diagrammatical shorthand, which is a counterpart of [10, (5.27)–(5.28)]: we define the *internal bubbles*

$$\begin{aligned} & \text{Diagram 14} \\ & := \frac{1}{2} \text{Diagram 15} - \frac{1}{2} \sum_{n=0}^{-\lambda} (-1)^{-\lambda-n} \text{Diagram 16} = \frac{1}{2} \text{Diagram 17} - \frac{1}{2} \left[ \text{Diagram 18} \right]_{u^{-1}} , \end{aligned} \tag{3.38}$$

(degree  $-2\lambda - 2$ )

$$\begin{aligned} & \text{Diagram 19} \\ & := \frac{1}{2} \text{Diagram 20} - \frac{1}{2} \sum_{n=0}^{\lambda} (-1)^{\lambda-n} \text{Diagram 21} = \frac{1}{2} \text{Diagram 22} - \frac{1}{2} \left[ \text{Diagram 23} \right]_{u^{-1}} \end{aligned} \tag{3.39}$$

(degree  $2\lambda - 2$ )

We also introduce internal bubbles on downward strings by taking the mates of (3.38) and (3.39), so that we maintain invariance under planar isotopy. It can be checked easily that internal bubbles commute with open dots and with other internal bubbles on the same string.

**Lemma 3.4.** *The following hold:*

- (1)  $\text{Diagram 24} = 2 \text{Diagram 25} - 2\delta_{\lambda,2} \text{Diagram 26}$  if  $\lambda \leq 2$ ;
- (2)  $\text{Diagram 27} = 2 \text{Diagram 28} - 2\delta_{\lambda,-2} \text{Diagram 29}$  if  $\lambda \geq -2$ .

*Proof.* We just prove (1), then (2) follows by applying the symmetry  $R$  (which interchanges the two sorts of internal bubble). Suppose that  $\lambda \leq 2$ . The idea is to compute

$$-\lambda \left[ \text{diagram with bubble on left} \right] = -\lambda \left[ \text{diagram with bubble on right} \right] \tag{3.40}$$

in two different ways. Starting from the form on the left-hand side, we slide the dot on the bubble past the crossing above it using (3.37) and the first relation from (3.6) to obtain

$$-\lambda \left[ \text{diagram with dot on bubble} \right] = -\lambda \left[ \text{diagram with dot on line} \right]$$

Then we slide the dot on the curl up past the crossing using (3.37) again to get

$$\begin{aligned} \lambda \left[ \text{diagram 1} \right] - \lambda \left[ \text{diagram 2} \right] &\stackrel{(3.23)}{=} \frac{1}{2} \lambda \left[ \text{diagram 3} \right] - \frac{1}{2} \sum_{n=0}^{\lambda} \lambda \left( \overset{\circ}{n} \right) \left[ \text{diagram 4} \right] \\ &\stackrel{(3.39)}{=} \lambda \left[ \text{diagram 5} \right] - \delta_{\lambda,1} \lambda \left[ \text{diagram 6} \right] - \delta_{\lambda,2} \lambda \left( \overset{\circ}{1} \right) \left[ \text{diagram 7} \right]. \end{aligned}$$

On the other hand, we can simplify the right-hand side of (3.40) using the second relation from (3.8) to obtain

$$\begin{aligned} 2 \lambda \left[ \text{diagram 8} \right] - \sum_{n=0}^{\lambda-1} \sum_{\substack{r,s \geq 0 \\ r+s=\lambda-1-n}} \lambda \left( \overset{\circ}{n} \right) \left[ \text{diagram 9} \right] \\ = 2 \lambda \left[ \text{diagram 10} \right] - \delta_{\lambda,1} \lambda \left[ \text{diagram 11} \right] - 2\delta_{\lambda,2} \lambda \left[ \text{diagram 12} \right] - \delta_{\lambda,2} \lambda \left( \overset{\circ}{1} \right) \left[ \text{diagram 13} \right]. \end{aligned}$$

The desired equality follows. □

**Lemma 3.5.** *The clockwise and counterclockwise internal bubbles are the two-sided inverses of each other, i.e., we have that*

$$\left[ \text{diagram 14} \right] \lambda = \left[ \text{diagram 15} \right] = \left[ \text{diagram 16} \right] \lambda .$$

*Proof.* We prove the second equality in the case that  $\lambda \geq -1$ . The first equality in this case then follows since internal bubbles commute, and after that the result may be deduced when  $\lambda \leq -1$  too by applying the symmetry  $R$ . So assume that  $\lambda \geq -1$ . By Lemma 3.4 (2) and the definition (3.39) of the counterclockwise internal bubble, we have that

$$\left[ \text{diagram 17} \right] \lambda = \left[ \text{diagram 18} \right] \lambda - \left[ \text{diagram 19} \right]_{u^{-1}} .$$



Now we observe that

$$\begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \stackrel{(3.9)}{=} \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} + \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \stackrel{(3.37)}{=} \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \stackrel{(3.23)}{=} \left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}}.$$

Substituting this into the previous formula gives

$$\begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} = \left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} - \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}} \stackrel{(3.34)}{=} \left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}} = \begin{array}{c} \uparrow \\ \lambda \end{array}.$$

To get the last equality here, we used that  $\left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}} = 1_{1_{\lambda+2}}$ , which follows from (3.21) on expanding the definitions (3.19) and (3.20).  $\square$

**Lemma 3.6.**  $\begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} + \begin{array}{c} \circlearrowright \\ \circlearrowright \\ \uparrow \\ \lambda \end{array} = \frac{1-(-1)^\lambda}{2} 1_{1_\lambda}.$

*Proof.* Expanding the definitions of the internal bubbles, also multiplying both sides by 2 and using the identity  $\begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} + \begin{array}{c} \circlearrowright \\ \circlearrowright \\ \uparrow \\ \lambda \end{array} = \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array}$  which follows by (3.33), this reduces to checking that

$$\begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} - \sum_{n=0}^{\lambda} (-1)^{\lambda-n} \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} - \sum_{n=0}^{-\lambda} (-1)^{-\lambda-n} \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} = (1 - (-1)^\lambda) 1_{1_\lambda}.$$

This follows in the three cases  $\lambda > 0$ ,  $\lambda = 0$  or  $\lambda < 0$  using (3.10) and Lemma 3.2.  $\square$

**Lemma 3.7.** *We have that*

$$\begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} - \left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}} = \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} - \left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}}$$

for either choice of orientation of the left-hand string.

*Proof.* We are going to expand

$$- \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array}$$

in two different ways, neither of which depends on the chosen orientation of the left-hand string. Commuting the dot on the bubble past the crossing above it using (3.37) and the first relation from (3.6) gives

$$\begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \stackrel{(3.37)}{=} \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} - \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \stackrel{(3.23)}{=} \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} - \left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}} \\ \stackrel{(3.34)}{=} \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} - \left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}} - \left[ \begin{array}{c} \circlearrowleft \\ \circlearrowleft \\ \uparrow \\ \lambda \end{array} \right]_{u^{-1}}.$$

On the other hand, we can apply (3.24) to obtain

$$\begin{aligned}
 & \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{u^{-1}} \\
 & \stackrel{(3.34)}{=} \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{u^{-1}} \\
 & \stackrel{(3.26)}{=} \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{u^{-1}} .
 \end{aligned}$$

The last terms produced in each case cancel with each other, and the Lemma 3.7 is proved.  $\square$

**Corollary 3.8.**  $\left[ \text{diagram} \right]_{\lambda} = \frac{1}{2} \left[ \text{diagram} \right]_{\lambda} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}}$

*Proof.* This follows from Lemma 3.7 (taking the left-hand string to be oriented downward) using also the definition of the internal bubble that is the mate of (3.38).  $\square$

**Corollary 3.9.**  $\left[ \text{diagram} \right]_{\lambda} = 2 \left[ \text{diagram} \right]_{\lambda}$  assuming  $\lambda \leq -2$ .

*Proof.* This follows from Lemma 3.7 (taking the left-hand string to be oriented upward) using also (3.18) and Lemma 3.4(1).  $\square$

**Lemma 3.10.**  $\left[ \text{diagram} \right]_{\lambda} = \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{u^{-1}}$

*Proof.* We have that

$$\begin{aligned}
 & \left[ \text{diagram} \right]_{\lambda} \stackrel{(3.24)}{=} \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{u^{-1}} \stackrel{(3.37)}{=} \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{u^{-1}} \\
 & \stackrel{(3.37)}{=} \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{u^{-1}} \\
 & \stackrel{(3.22)}{=} \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{u^{-1}} .
 \end{aligned}$$

$\square$

**Lemma 3.11.**  $\left[ \text{diagram} \right]_{\lambda} = \left[ \text{diagram} \right]_{\lambda}$  and  $\left[ \text{diagram} \right]_{\lambda} = \left[ \text{diagram} \right]_{\lambda}$ .

*Proof.* We prove the first equality assuming that  $\lambda \leq -2$ . Then the second equality for  $\lambda \leq -2$  can be deduced using Lemma 3.5, and after that both equalities for  $\lambda \geq -2$  follow by applying R. So assume that  $\lambda \leq -2$ . By Lemma 3.4(1) and Corollary 3.9, we have that

We need to show that this commutes with the crossing  $\times^{\lambda}$ . It is easy to check that the various dots do so, indeed,  $x_1x_2$  and  $x_1 + x_2$  are central elements in the nil-Hecke algebra  $NH_2$ . It is almost as easy to see that the counterclockwise bubbles commute with the crossing too:

The Lemma 3.11 is proved. □


**Lemma 3.12.**


*Proof.* We just have to use the rotations of (3.35) then (3.36) to commute dots downward past the crossing:

**Corollary 3.13.** □



*Proof.* In the identity from Lemma 3.12, we use the image of Corollary 3.8 under  $\mathbb{R}$  to combine the first and fourth terms on the right-hand side to obtain the identity

It remains to rotate clockwise by  $90^\circ$  then apply R. □

**Lemma 3.14.**   $\lambda = \frac{1}{2} \downarrow_{-1} \lambda - \downarrow_{-1} \lambda.$

*Proof.* Closing the identity established in Lemma 3.12 on the top with a rightward cap and using (3.22) to expand the two curls produced gives that  where

$$\begin{aligned}
 A &= -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} \stackrel{(3.34)}{=} -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} \\
 &\stackrel{(3.33)}{=} \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}}, \\
 B &= \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} \stackrel{(3.33)}{=} -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} + \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}}, \\
 C &= \text{diagram}, \\
 D &= \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}, v^{-1}} \stackrel{(3.18)}{=} \delta_{\lambda, 0} \frac{1}{2} \downarrow_{-1} \lambda, \\
 E &= -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}}.
 \end{aligned}$$


After rewriting  $A$ ,  $B$  and  $D$  as indicated, the first terms from the expansions of  $A$  and  $B$  combine according to the definition (3.38) to produce the second term  from the formula we are trying to prove. Similarly, the second terms from the expansions of  $A$  and  $B$  combine to give . Adding this to  $C$  and  $D$  and applying Lemma 3.6 gives  $\frac{1-(-1)^\lambda + \delta_{\lambda, 0}}{2} \downarrow_{-1} \lambda$ . Comparing with the first term in the formula we are trying to prove, we are left with showing that the remaining term from  $A$  plus the term from  $E$  equals  $\frac{(-1)^\lambda - \delta_{\lambda, 0}}{2} \downarrow_{-1} \lambda$ . This holds because these two terms can be simplified as follows:

$$-\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} = \begin{cases} \frac{(-1)^\lambda}{2} \downarrow_{-1} \lambda & \text{if } \lambda < 0 \\ 0 & \text{otherwise,} \end{cases} \tag{3.41}$$


$$-\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} = \begin{cases} \frac{(-1)^\lambda}{2} \downarrow_{-1} \lambda & \text{if } \lambda > 0 \\ 0 & \text{otherwise.} \end{cases} \tag{3.42}$$

The arguments to prove (3.41) and (3.42) are similar, so we just explain how to see the second one. The counterclockwise fake bubble polynomial is 0 if  $\lambda < 0$ , so the expression on the left-hand side of (3.42) is 0 in this case, as required. The fake bubble polynomial is the identity if  $\lambda = 0$ , and it is easy to see that (3.42) is true in this case too due to


the presence of two pins. Finally, if  $\lambda > 0$ , the fake bubble polynomial is  $u^\lambda +$  (lower terms), and by (3.10) we have that  $\textcircled{u}^\pm \textcircled{\circ}^\lambda = (-1)^{\lambda-1} u^{-\lambda} +$  (lower terms). It follows that the  $u^{-1}$ -coefficient on the left-hand side of (3.42) comes from the leading terms, and the identity follows easily.  $\square$

**Corollary 3.15.** 

*Proof.* By a rotated version of (3.37), we have that



It remains to apply Lemma 3.14 to the first term on the right-hand side.  $\square$

**Lemma 3.16.** 

*Proof.* We begin by attaching a rightward crossing to the top of the identity from Lemma 3.12 to get that  $\textcircled{\circ}^\lambda = A + B + C + D + E$  where

$$\begin{aligned}
 A &= \frac{1}{2} \textcircled{\circ}^\lambda \stackrel{(3.24)}{=} -\frac{1}{2} \textcircled{\circ}^\lambda + \frac{1}{2} \left[ \textcircled{u}^\pm \textcircled{\circ}^\lambda \right]_{u^{-1}} \\
 &\stackrel{(3.34)}{=} -\frac{1}{2} \textcircled{\circ}^\lambda + \frac{1}{2} \left[ \textcircled{u}^\pm \textcircled{\circ}^\lambda \right]_{u^{-1}} + \frac{1}{2} \left[ \textcircled{u}^\pm \textcircled{\circ}^\lambda \right]_{u^{-1}} \\
 &\stackrel{(3.33)}{=} -\frac{1}{2} \textcircled{\circ}^\lambda - \frac{1}{2} \left[ \textcircled{u}^\pm \textcircled{\circ}^\lambda \right]_{u^{-1}} + \frac{1}{2} \left[ \textcircled{u}^\pm \textcircled{\circ}^\lambda \right]_{u^{-1}} \\
 &\quad + \frac{1}{2} \left[ \textcircled{u}^\pm \textcircled{\circ}^\lambda \right]_{u^{-1}},
 \end{aligned}$$

$$\begin{aligned}
 B &= \frac{1}{2} \text{diagram} \\
 &\stackrel{(3.37)}{=} \frac{1}{2} \text{diagram} + \frac{1}{2} \text{diagram} \stackrel{(3.22)}{=} -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} + \frac{1}{2} \text{diagram} \\
 &\stackrel{(3.33)}{=} -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} \\
 &\stackrel{(3.34)}{=} -\frac{1}{2} \text{diagram} - \frac{1}{2} \text{diagram} \\
 &\stackrel{(3.33)}{=} \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} - \text{diagram} + \text{diagram}, \\
 C &= \text{diagram} = \frac{1}{2} \text{diagram} - \text{diagram} \quad (\text{by Lemma 3.14 and duality}), \\
 D &= -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} \stackrel{(3.24)}{=} \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}, v^{-1}}, \\
 E &= -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} \stackrel{(3.36)}{=} -\frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}} \\
 &\stackrel{(3.22)}{=} \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}, v^{-1}} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}}.
 \end{aligned}$$

It just remains to gather the terms together to see that we get exactly the three terms on the right-hand side of the formula claimed in the statement of the Lemma 3.16. The first terms from  $A$  and  $D$  combine according to Corollary 3.8 (rotated through  $180^\circ$ ) to give the second term in the claimed formula. The second term from  $A$  cancels with the first term from  $B$ . The third term from  $B$  gives the third term in the claimed formula. The fourth term from  $B$  cancels with the second term from  $C$ . The first term from  $C$  gives the first term in the claimed formula. This just leaves the second term in  $B$ , the second term in  $D$  and both terms in  $E$ . The second term in  $D$  and the first term in  $E$  are easily seen to equal 0 using the definitions (3.18). The second terms in  $B$  and  $E$  are 0 too, as follows by considering leading terms like in the proof of (3.41) and (3.42).  $\square$

**Lemma 3.17.** 
$$\left[ \text{diagram} \right]_{u^{-1}} = \text{diagram} + \text{diagram} - \frac{1}{2} \left[ \text{diagram} \right]_{u^{-1}}.$$

*Proof.* We begin by calculating:

$$\begin{aligned}
 & \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \stackrel{(3.37)}{=} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \stackrel{(3.37)}{=} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \\
 & \stackrel{(3.22)}{=} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \\
 & \stackrel{(3.23)}{=} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \\
 & \stackrel{(3.34)}{=} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \\
 & \quad + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \\
 & \stackrel{(3.34)}{=} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \\
 & \quad + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda .
 \end{aligned}$$

In deriving the last equality here, we have removed two terms that are 0 by the same argument as used to prove (3.41) and (3.42). Then we use (3.39) to combine the first and third terms to obtain the identity

$$\begin{aligned}
 & \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \\
 & = -2 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda .
 \end{aligned}$$

Similarly, we use Corollary 3.8 (or rather its image under  $\tau$ ) to combine the term on the left-hand side with the fourth term on the right-hand side to reduce to

$$2 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda = -2 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda .$$

Dividing by 2 and adding an inverse dot to the right-hand string, this rearranges to show that

$$\begin{aligned}
 & \frac{1}{2} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \frac{1}{2} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda \\
 & = \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \frac{1}{2} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda ,
 \end{aligned}$$

the additional second term on the left-hand side obviously being equal to 0. It remains to observe that the left-hand side in this is equal to the left-hand in the formula we are trying to prove by (3.39).  $\square$

**Lemma 3.18.**  $\left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda = \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + 2 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda + 2 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda - \delta_{\lambda, -1} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \left| \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right. \right] \lambda$  for  $\lambda \leq -1$ .

*Proof.* We expand the left-hand side:

$$\begin{aligned}
 & \left[ \text{diagram} \right]_{\lambda} \stackrel{(3.24)}{=} \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{\lambda} \stackrel{(3.37)}{=} \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{\lambda} - \left[ \text{diagram} \right]_{\lambda} \\
 & \stackrel{(3.37)}{=} \left[ \text{diagram} \right]_{u^{-1}} + \left[ \text{diagram} \right]_{\lambda} - \left[ \text{diagram} \right]_{\lambda} - \left[ \text{diagram} \right]_{\lambda} \\
 & \stackrel{(3.37)}{=} \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{\lambda} - \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} \\
 & \stackrel{(3.37)}{=} \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{\lambda} - \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} \\
 & \stackrel{(3.23)}{=} \left[ \text{diagram} \right]_{u^{-1}} + \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda}.
 \end{aligned}$$

Now we use the definition (3.39) to rewrite the fourth and sixth terms in this expression:

$$\begin{aligned}
 & \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} = 2 \left[ \text{diagram} \right]_{\lambda} + 2 \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{u^{-1}} + \left[ \text{diagram} \right]_{u^{-1}} \\
 & \stackrel{(3.34)}{=} 2 \left[ \text{diagram} \right]_{\lambda} + 2 \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} \\
 & \quad - \left[ \text{diagram} \right]_{\lambda} + \left[ \text{diagram} \right]_{\lambda} - \left[ \text{diagram} \right]_{\lambda}.
 \end{aligned}$$

It just remains to expand the fake bubble polynomials using the assumption that  $\lambda \leq -1$  to see that

$$\left[ \text{diagram} \right]_{\lambda} - \left[ \text{diagram} \right]_{\lambda} - \left[ \text{diagram} \right]_{\lambda} = -\delta_{\lambda, -1} \left[ \text{diagram} \right]_{\lambda}.$$

The Lemma 3.18 is proved. □

#### 4. MONOIDAL FUNCTOR FROM $\mathcal{N}\mathcal{B}_t$ TO A LOCALIZED VERSION OF $\mathfrak{U}(\mathfrak{sl}_2)$

We are now in position to construct the strict monoidal functor

$$\Omega_t : \mathcal{N}\mathcal{B}_t \rightarrow \text{Add} \left( \mathfrak{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ} \right).$$

The latter category is the additive envelope of a monoidal category obtained by collapsing the 2-categorical structure on  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ \dots \circ}$ . Its full definition is as follows:

**Definition 4.1.** Let  $\mathfrak{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ}$  be the monoidal category with objects that are words in the free monoid  $\langle E, F \rangle$  generated by the letters  $E$  and  $F$ . For any  $X, Y \in \langle E, F \rangle$  and  $\lambda \in \mathbb{Z}$ , there are corresponding 1-morphisms  $X1_{\lambda}, Y1_{\lambda}$  in  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ \dots \circ}$  obtained by horizontally composing the 1-morphisms  $E1_{\mu}$  and  $F1_{\mu}$  corresponding to the letters of  $X$  and  $Y$  for appropriate weights  $\mu$ . Then we define

$$\text{Hom}_{\mathfrak{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ}}(Y, X) := \prod_{\lambda \in t+2\mathbb{Z}} \text{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)_{\circ \dots \circ}}(Y1_{\lambda}, X1_{\lambda}) \tag{4.1}$$



for  $X, Y \in \langle E, F \rangle$ . Defining the *weight*  $\text{wt}(X)$  of  $X \in \langle E, F \rangle$  to be  $2 \times$  (the number of letters  $E$  minus the number of letters  $F$  in the word  $X$ ), the morphism space (4.1) is 0 unless  $\text{wt}(X) = \text{wt}(Y)$ . In general,  $f \in \text{Hom}_{\mathcal{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ}}(Y, X)$  is a tuple  $f = (f_\lambda)_{\lambda \in 1+2\mathbb{Z}}$  of morphisms  $f_\lambda \in \text{Hom}_{\mathcal{U}_{q^{-1}(\mathfrak{sl}_2)_{\circ \dots \circ}}}(Y1_\lambda, X1_\lambda)_{\circ \dots \circ}$ . The composition law making  $\mathcal{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ}$  into a category is induced by vertical composition in  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ \dots \circ}$ : we have that  $(g \circ f)_\lambda = g_\lambda \circ f_\lambda$  for morphisms  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ . The strict monoidal product  $- \star - : \mathcal{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ} \boxtimes \mathcal{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ} \rightarrow \mathcal{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ}$  is induced by horizontal composition in  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ \dots \circ}$ : it is defined on objects simply by concatenation of words and on morphisms by setting  $(f' \star f)_\lambda := f'_{\lambda + \text{wt}(Y)} f_\lambda$  for  $f : Y \rightarrow X$ ,  $f' : Y' \rightarrow X'$ .

Morphisms in the additive envelope  $\text{Add}(\mathcal{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ})$  are matrices of morphisms in  $\mathcal{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ}$ . In the statement of the next theorem, we use some obvious shorthand to represent such matrices. For example, the morphism  $\Omega_t(\times)$  appearing below is an endomorphism of

$$(E \oplus F)^{\star 2} \cong E \star E \oplus E \star F \oplus F \star E \oplus F \star F,$$

so it is a  $4 \times 4$  matrix with rows and columns indexed by the words  $EE, EF, FE$  and  $FF$ . In turn,  $\Omega_t(\times)_\lambda$  is a matrix representing an endomorphism of  $EE1_\lambda \oplus \widehat{E}F1_\lambda \oplus FE1_\lambda \oplus FF1_\lambda$ . The eight morphisms appearing on the right-hand side of the equation for  $\Omega_t(\times)_\lambda$  in the statement of the theorem are matrices of this form with 0 in all but the self-evident entry.

**Theorem 4.2.** *There is a strict monoidal functor  $\Omega_t : \mathfrak{NB}_t \rightarrow \text{Add}(\mathcal{U}(\mathfrak{sl}_2; t)_{\circ \dots \circ})$  taking the generating object  $B$  to  $E \oplus F$ , and defined on generating morphisms by letting*

$$\begin{aligned} \Omega_t \left( \begin{array}{c} \downarrow \\ \circ \\ \downarrow \end{array} \right)_\lambda &:= \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \lambda - \begin{array}{c} \downarrow \\ \circ \\ \downarrow \end{array} \lambda, \\ \Omega_t \left( \times \right)_\lambda &:= \begin{array}{c} \nearrow \\ \lambda \\ \searrow \end{array} + \begin{array}{c} \nwarrow \\ \lambda \\ \nearrow \end{array} + \begin{array}{c} \nearrow \\ \lambda \\ \nearrow \end{array} + \begin{array}{c} \nearrow \\ \lambda \\ \searrow \end{array} + \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \lambda \\ &\quad - \begin{array}{c} \nearrow \\ \lambda \\ \searrow \end{array} + \begin{array}{c} \nwarrow \\ \lambda \\ \nearrow \end{array} + \begin{array}{c} \nearrow \\ \lambda \\ \nearrow \end{array}, \\ \Omega_t \left( \cap \right)_\lambda &:= \begin{array}{c} \curvearrowright \\ \lambda \end{array} + \begin{array}{c} \curvearrowright \\ \lambda \end{array}, \\ \Omega_t \left( \cup \right)_\lambda &:= \begin{array}{c} \curvearrowleft \\ \lambda \end{array} + \begin{array}{c} \curvearrowleft \\ \lambda \end{array} \end{aligned}$$

for  $\lambda \in t + 2\mathbb{Z}$ . We also have that

$$\Omega_t(\mathbb{O}(u))_\lambda = \lambda \begin{array}{c} \circ \\ \downarrow \end{array} (-u) \begin{array}{c} \circ \\ \downarrow \end{array} (u). \tag{4.2}$$

*Proof.* To prove the existence of  $\Omega_t$ , we simply need to check the eight defining relations from (2.2) to (2.5)!

- Consider the first relation from (2.3). We have that

$$\Omega_t \left( \bigcirc \right)_\lambda = \begin{array}{c} \curvearrowright \\ \lambda \end{array} + \begin{array}{c} \curvearrowleft \\ \lambda \end{array}.$$

To check the relation, we must show that this equals  $\Omega_t(t1_\mathbb{1})_\lambda = t1_{1_\lambda}$ , which follows immediately from Lemma 3.6.

- For the second relation from (2.3), we have that

$$\begin{aligned} \Omega_t \left( \bigcup \right)_{\lambda} &= \bigcup_{\lambda} + \text{bubble} \lambda, \\ \Omega_t \left( \big| \right)_{\lambda} &= \uparrow_{\lambda} + \downarrow_{\lambda}, \\ \Omega_t \left( \bigcup \right)_{\lambda} &= \text{dot-bubble} \lambda + \bigcup_{\lambda}. \end{aligned}$$

Thus, to check this relation, we need to show that

$$\bigcup_{\lambda} = \uparrow_{\lambda} = \text{dot-bubble} \lambda, \quad \text{bubble} \lambda = \downarrow_{\lambda} = \bigcup_{\lambda}.$$

These both follow easily using (3.7), (3.13) and Lemma 3.5.

- In a similar way, the first relation from (2.5) reduces to checking that

$$\begin{aligned} \text{dot-cross} \lambda - \text{cross-dot} \lambda &= \uparrow \downarrow \lambda, & -\text{dot-cross} \lambda + \text{cross-dot} \lambda &= \downarrow \uparrow \lambda, \\ -\text{dot-cross} \lambda + \text{cross-dot} \lambda &= -\text{cup} \lambda, & \text{dot-bubble-cross} \lambda - \text{cross-dot-bubble} \lambda &= -\text{dot-bubble} \lambda, \end{aligned}$$

and that

$$\begin{aligned} \text{dot-dot} \lambda - \text{dot-bubble} \lambda + \text{dot-dot} \lambda - \text{dot-bubble} \lambda &= \uparrow \downarrow \lambda - \text{cup} \lambda, \\ \text{dot-dot} \lambda - \text{dot-bubble} \lambda + \text{dot-dot} \lambda - \text{dot-bubble} \lambda &= \uparrow \downarrow \lambda - \text{cup} \lambda. \end{aligned}$$

The first four of these follow immediately from the last defining relation in (3.6) plus its variants obtained by rotating through 90°, 180° and 270°. The last two follow using (3.30).

- Next, we look at the second relation from (2.5). For this, we must show that

$$\text{dot-cup} \lambda = \text{cup-dot} \lambda, \quad \text{dot-bubble} \lambda = \text{dot-bubble} \lambda.$$

These both follow using planar isotopy since open dots commute with internal bubbles.

- The first relation in (2.4) follows using Lemma 3.5 and the relations

$$\begin{aligned} \text{dot-bubble} \lambda + \text{dot-bubble} \lambda - \frac{1}{2} \text{dot-cup} \lambda &= 0, \\ \text{dot-bubble} \lambda - \text{dot-bubble} \lambda + \frac{1}{2} \text{dot-cup} \lambda &= 0. \end{aligned}$$


The first of these is Lemma 3.14, and the second follows from the first (with  $\lambda$  replaced by  $-\lambda$ ) on applying R.

- The second relation from (2.4) requires the following four identities to be checked:




The first two of these follow immediately using planar isotopy. The last one follows using planar isotopy and Lemma 3.5. For the third, one also needs to use Lemma 3.11 (rotated counterclockwise by 90°) to commute the two clockwise internal bubbles past the crossing to their left.

- The first relation from (2.2) involves six non-trivial relations, coming from the  $(EE1_\lambda, EE1_\lambda)$ -  $(EF1_\lambda, FE1_\lambda)$ -,  $(FE1_\lambda, FE1_\lambda)$ -,  $(FF1_\lambda, FF1_\lambda)$ -,  $(FE1_\lambda, EF1_\lambda)$ - and  $(EF1_\lambda, EF1_\lambda)$ -entries of the corresponding  $4 \times 4$  matrices. However, after applying Lemma 3.5 to redistribute some internal bubbles, the last three may be deduced from images of the first three under  $\mathbb{R}$  and  $\mathbb{T}$ . Thus, we really only have to verify three relations, which are as follows:



The first of these is the first defining relation from (3.6). The third one follows from Lemma 3.16 after making an obvious application of (3.33). To prove the second one, we take the equation from Corollary 3.15 and add it to the equation obtained from it by applying  $\mathbb{T}$  to both sides to deduce that



Now use Lemma 3.3.

- The second relation from (2.2) is the most complicated to check since it involves an equality of  $8 \times 8$  matrices, and there are 20 non-zero entries in these matrices. After simplifying with Lemmas 3.5, 3.11 and using the symmetries  $\mathbb{R}$  and  $\mathbb{T}$ , the calculation reduces to checking five relations, coming from the  $(EEE1_\lambda, EEE1_\lambda)$ -,  $(FEE1_\lambda, EEF1_\lambda)$ -,  $(EEF1_\lambda, EEF1_\lambda)$ -,  $(EFE1_\lambda, EEF1_\lambda)$ - and  $(EFE1_\lambda, EFE1_\lambda)$ -entries. The first two of these are



both of which follow from the second defining relation in 3.7 (using also some planar isotopy to get the one on the right). The relations from the  $(EEF1_\lambda, EEF1_\lambda)$ - and  $(EFE1_\lambda, EEF1_\lambda)$ -entries are

We leave the verification of these as exercises for the reader—simplify using Corollary 3.15 and Lemma 3.16.

Finally, we must verify the relation arising from the  $(EFE1_\lambda, EFE1_\lambda)$ -entry. We found this to be significantly harder than the relations encountered so far! After rearranging terms and simplifying a little using Lemmas 3.5 and 3.11, it requires the following to be proved:

Notice that the right-hand side is the image of the left-hand side with  $\lambda$  replaced by  $-\lambda - 2$  under the map  $-R$ . Thus, we must show that  $A_\lambda + R(A_{-\lambda-2}) = 0$  where  $A_\lambda$  is the expression consisting of the nine terms on the left-hand side of the relation. We show equivalently that  $B_\lambda + R(B_{-\lambda-2}) = 0$  where

To obtain the second expression for  $B_\lambda$ , we have used 3.30 several times to split the fourth term and the last term into two and to combine the fifth, sixth, seventh and eighth terms into one. We denote the eight terms in this by  $B_{\lambda;1}, \dots, B_{\lambda;8}$  in order from left to right, so  $B_\lambda = B_{\lambda;1} + \dots + B_{\lambda;8}$ . By Corollary 3.13, we have that  $B_{\lambda;1} = C_{\lambda;1} + C_{\lambda;2} + C_{\lambda;3} + C_{\lambda;4}$  where

$$\begin{aligned}
 C_{\lambda;1} &= \text{diagram} \stackrel{(3.6)}{=} \text{diagram} - \text{diagram} - \text{diagram} \\
 &= \text{diagram} + \text{diagram} - 1 \circ \text{diagram} - \text{diagram} \quad (\text{by Lemma 3.3}) \\
 &\stackrel{(3.24)}{=} \text{diagram} - \text{diagram} + \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} + \text{diagram} \\
 &\quad - \frac{1}{2} \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} + \text{diagram} - \frac{1}{2} \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} \\
 &= \text{diagram} + \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} \\
 &\quad - \frac{1}{2} \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} - \frac{1}{2} \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} - \text{diagram} - B_{\lambda;2} - B_{\lambda;3},
 \end{aligned}$$

$$\begin{aligned}
 C_{\lambda;2} &= -2 \text{diagram} = 2 \text{diagram} + 4 \text{diagram} \\
 &= -B_{\lambda;5} - B_{\lambda;6} - B_{\lambda;7} \quad (\text{by Lemma 3.16}),
 \end{aligned}$$

$$\begin{aligned}
 C_{\lambda;3} &= - \text{diagram} \stackrel{(3.37)}{=} - \text{diagram} + \text{diagram} \\
 &\stackrel{(3.24)}{=} 2 \text{diagram} - \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} + \text{diagram} - \text{diagram} \\
 &\stackrel{(3.23)}{=} 2 \text{diagram} - \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} + \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} - \text{diagram} \\
 &\stackrel{(3.34)}{=} 2 \text{diagram} - \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} + \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} - \text{diagram} \\
 &\stackrel{(3.39)}{=} 2 \text{diagram} - \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} - 2 \text{diagram} \\
 &\stackrel{(3.34)}{=} 2 \text{diagram} - \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} + \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} - B_{\lambda;4}, \\
 &= 2 \text{diagram} - \frac{1}{2} \left[ \begin{array}{c} u \\ \text{diagram} \\ u \end{array} \right]_{u^{-1}} - B_{\lambda;4},
 \end{aligned}$$

$$\begin{aligned}
 C_{\lambda;4} &= \left[ \text{diagram} \right]_{u^{-1}} \\
 &\stackrel{(3.36)}{=} \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{u^{-1}} \\
 &\stackrel{(3.24)}{=} -2 \left[ \text{diagram} \right]_{u^{-1}} + \left[ \text{diagram} \right]_{u^{-1},v^{-1}} \\
 &\stackrel{(3.36)}{=} -2 \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{u^{-1},v^{-1}} + \left[ \text{diagram} \right]_{u^{-1}} \\
 &\stackrel{(3.23)}{=} -2 \left[ \text{diagram} \right]_{u^{-1}} - \left[ \text{diagram} \right]_{u^{-1},v^{-1}} = -2 \left[ \text{diagram} \right]_{u^{-1}}.
 \end{aligned}$$

For the final equality in the expansion of  $C_{\lambda;3}$ , the third term on the previous line is 0 by a similar argument to the proof of (3.41) and (3.42). To see that the second terms on the two lines are equal, this is clear when  $\lambda < -2$  for the simple reason that the counterclockwise fake bubble polynomial is 0 in that case, and it is almost as easy to see that both terms are 0 when  $\lambda = -2$ . So we may assume that  $\lambda > -2$ . Then the clockwise bubble can be converted into an internal bubble using Lemma 3.4(2), after which it cancels with the counterclockwise internal bubble by Lemma 3.5. In the expansion of  $C_{\lambda;4}$ , three terms have been removed without explicit reference since they are 0 by the usual arguments. Now we collect the pieces to see that  $B_\lambda = D_\lambda + E_\lambda + F_\lambda$  where

$$\begin{aligned}
 D_\lambda &= \left( \text{diagram} \right)_\lambda - \left[ \text{diagram} \right]_{u^{-1}} \quad (\text{the second term is new, it cancels a term on the next line}), \\
 E_\lambda &= \frac{1}{2} \left[ 2 \left( \text{diagram} \right)_\lambda + 2 \left( \text{diagram} \right)_{-1} \lambda - \left( \text{diagram} \right)_{-1} \lambda - \left( \text{diagram} \right)_{-1} \lambda - \left( \text{diagram} \right)_{-1} \lambda \right]_{u^{-1}}, \\
 F_\lambda &= 4 \left( \text{diagram} \right)_\lambda - \left( \text{diagram} \right)_\lambda + 2 \left( \text{diagram} \right)_\lambda - 2 \left[ \text{diagram} \right]_{u^{-1}}. \tag{4.3}
 \end{aligned}$$

Next, observe that  $D_\lambda + R(D_{-\lambda-2}) = 0$  by (3.27). Also  $E_\lambda = 0$ , indeed, already the expression inside the square brackets vanishes as follows from the elementary identity

$$\frac{2}{(u-x)(u-y)(u-z)} + \frac{2}{(u-x)(u-y)z} - \frac{1}{(u-y)(u-z)z} - \frac{1}{(u-x)(u-z)z} - \frac{x+y}{(u-x)(u-y)(u-z)z} = 0$$

where  $x$  represents the dot on the top left component,  $y$  the dot on the bottom left component, and  $z$  the dot on the rightmost vertical string.

In view of all of this, it just remains to show that  $F_\lambda + R(F_{-\lambda-2}) = 0$ . Since  $\tau$  is an involution and we always have that  $\lambda \leq -1$  or  $-\lambda - 2 \leq -1$ , we may assume without loss of generality that  $\lambda \leq -1$ . Hence, we have that

$$\left( \text{diagram} \right)_\lambda = 0, \quad \left( \text{diagram} \right)_\lambda = \begin{cases} u \uparrow \downarrow \uparrow \lambda + \uparrow \downarrow \textcircled{1} \uparrow \lambda & \text{if } \lambda = -1 \\ \uparrow \downarrow \uparrow \lambda & \text{if } \lambda = -2 \\ 0 & \text{if } \lambda \leq -3. \end{cases} \tag{4.4}$$

From (4.3) and using the first of the equalities (4.4), we have that

$$\begin{aligned}
 R(F_{-\lambda-2}) &= 4 \left[ \text{diagram} \right] - \left[ \text{diagram} \right] \left| \lambda + 2 \right. \left[ \text{diagram} \right] \\
 &\stackrel{(3.30)}{=} -4 \left[ \text{diagram} \right] + 4 \left[ \text{diagram} \right] - \left[ \text{diagram} \right] \left| \lambda - 2 \right. \left[ \text{diagram} \right] + 2 \left[ \text{diagram} \right] \left| \lambda \right. \\
 &= -4 \left[ \text{diagram} \right] + 4 \left[ \text{diagram} \right] - \left[ \text{diagram} \right] \left| \lambda - 2 \right. \left[ \text{diagram} \right] + \left[ \text{diagram} \right] \left| \lambda \right. \\
 &\stackrel{(3.30)}{=} -4 \left[ \text{diagram} \right] + 4 \left[ \text{diagram} \right] + 2 \left[ \text{diagram} \right] \left| \lambda - 2 \right. \left[ \text{diagram} \right] \\
 &= -4 \left[ \text{diagram} \right] + 2 \left[ \text{diagram} \right],
 \end{aligned}$$

where we also used Lemma 3.4(1), Lemma 3.5 and Corollary 3.8 to get the last three equalities. Also by Lemma 3.17, we have that

$$\begin{aligned}
 F\lambda &= 4 \left[ \text{diagram} \right] - \left[ \text{diagram} \right] \left| \lambda - 2 \right. \left[ \text{diagram} \right] + \left[ \text{diagram} \right] \Big|_{u^{-1}} \\
 &\stackrel{(3.30)}{=} -4 \left[ \text{diagram} \right] + 4 \left[ \text{diagram} \right] - \left[ \text{diagram} \right] \left| \lambda + 2 \right. \left[ \text{diagram} \right] \\
 &\qquad\qquad\qquad - 2 \left[ \text{diagram} \right] \left| \lambda \right. + \left[ \text{diagram} \right] \Big|_{u^{-1}} \\
 &\stackrel{(3.39)}{=} -4 \left[ \text{diagram} \right] - \left[ \text{diagram} \right] \left| \lambda + 2 \right. \left[ \text{diagram} \right] - 2 \left[ \text{diagram} \right] \Big|_{u^{-1}} + \left[ \text{diagram} \right] \Big|_{u^{-1}} \\
 &\stackrel{(3.30)}{=} -4 \left[ \text{diagram} \right] - \left[ \text{diagram} \right] \left| \lambda - 2 \right. \left[ \text{diagram} \right] + 2 \left[ \text{diagram} \right] \left| \lambda \right. \\
 &\qquad\qquad\qquad - 2 \left[ \text{diagram} \right] \Big|_{u^{-1}} + \left[ \text{diagram} \right] \Big|_{u^{-1}} \\
 &= -4 \left[ \text{diagram} \right] - 2 \left[ \text{diagram} \right] + 2\delta_{\lambda,-1} \left[ \text{diagram} \right].
 \end{aligned}$$

The last equality here needs further explanation. We used Lemma 3.10 to rewrite the fourth term on the penultimate line to deduce that the sum of the fourth, fifth and sixth terms on this line equal

$$\begin{aligned}
 2 \left[ \text{diagram} \right] \left| \lambda - 2 \right. \left[ \text{diagram} \right] - 2 \left[ \text{diagram} \right] \left| \lambda \right. \\
 - 2 \left[ \text{diagram} \right] \Big|_{u^{-1}} + \left[ \text{diagram} \right] \Big|_{u^{-1}}. \tag{4.5}
 \end{aligned}$$

Using also Lemma 3.4(1) and Lemma 3.5, the first term in (4.5) simplifies further to  $\uparrow \downarrow -1 \uparrow \lambda$ , hence, it cancels with the second term on the penultimate line above. When  $\lambda \leq -3$ , it is easy to see from (4.4) that the remaining four terms in (4.5) are all 0, and the desired equality follows. If  $\lambda = -2$ , the remaining terms are 0 again—this time two of them are non-zero but they cancel with each other. Finally, when  $\lambda = -1$  the remaining four terms simplify by a calculation using also Corollary 3.8 to obtain the final term in the formula we are trying to prove. We use the simplified formulae for  $F_\lambda$  and  $R(F_{-\lambda-2})$  now established to see that  $F_\lambda + R(F_{-\lambda-2}) = 0$ . Indeed, after cancelling the clockwise internal bubble and a dot from the leftmost string, the sum is equal to

$$2 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda - 2 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda - 4 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda - 4 \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda + 2\delta_{\lambda,-1} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda,$$

which is 0 by Lemma 3.18.

It just remains to prove (4.2). By (3.19) and (3.20), it suffices to show that

$$(-1)^\lambda [\Omega_t(\mathbb{O}(u))]_{u < 0} = \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda.$$

Equivalently, since  $\left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda = - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda$ , we show that

$$(-1)^\lambda [\Omega_t(\mathbb{O}(u))]_{u < 0} = \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda.$$

By (2.22) and the definition of  $\Omega_t$ , we have that

$$(-1)^\lambda [\Omega_t(\mathbb{O}(u))]_{u < 0} = -2u \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda \Big|_{u < -1}.$$

Expanding the definitions of the internal bubbles using (3.38) and (3.39), this equals  $A + B + C$  where

$$\begin{aligned} A &= u \left[ \sum_{n=0}^{-\lambda} (-1)^{-\lambda-n} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda \right]_{u < -1}, \\ B &= u \left[ \sum_{n=0}^{\lambda} (-1)^{\lambda-n} \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda \right]_{u < -1}, \\ C &= -u \left[ \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda + \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda \right]_{u < -1}. \end{aligned}$$

We complete the proof by showing that

$$A = \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda, B = - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda \quad \text{and} \quad C = - \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \left[ \begin{array}{c} \uparrow \\ \circ \\ \downarrow \end{array} \right] \lambda.$$



For  $A$  and  $B$ , this follows simply by expanding the definitions of the pins and fake bubble polynomials, e.g., for  $B$  we have that

$$\begin{aligned}
 B &= u \sum_{n=0}^{\lambda} \sum_{r \geq 1} (-1)^{\lambda-n+r} \lambda_{-n-1+r} \circlearrowleft \circlearrowright \lambda u^{-r-1} \\
 &= - \sum_{n=0}^{\lambda} \sum_{r \geq \lambda-n} (-1)^r r \circlearrowleft \circlearrowright \lambda u^{\lambda-n-r-1} = - \left[ u \pm \circlearrowleft \circlearrowright \lambda \right]_{u < 0}.
 \end{aligned}$$

For  $C$ , we first use the relations (3.33) and (3.34) to see that

$$\circlearrowleft \circlearrowright \lambda + \circlearrowright \circlearrowleft \lambda = \circlearrowleft \circlearrowright \lambda + \circlearrowright \circlearrowleft \lambda.$$

Now we can expand the definitions to get that

$$\begin{aligned}
 C &= -u \sum_{r,s \geq 0} (-1)^r r \circlearrowleft \circlearrowright s-1 u^{-r-s-2} - u \sum_{r \geq 1} (-1)^r r-1 \circlearrowleft \circlearrowright -1 u^{-r-1} \\
 &= -u \sum_{\substack{r \geq 0 \\ s \geq 1}} (-1)^r r \circlearrowleft \circlearrowright s-1 u^{-r-s-2} \\
 &= - \sum_{r,s \geq 0} (-1)^r r \circlearrowleft \circlearrowright s u^{-r-s-2} = - \circlearrowleft \circlearrowright \lambda.
 \end{aligned}$$

This completes the proof of the Theorem 4.2. □

**Remark 4.3.** The monoidal functor in Theorem 4.2 is certainly not unique. One way to obtain alternative forms, maintaining the property (4.2) and preserving the leading terms  $\circlearrowleft \lambda$  and  $\circlearrowright \lambda$  in the formulae for  $\Omega_t(\bullet)_{\lambda}$  and  $\Omega_t(\times)_{\lambda}$ , is as follows. Suppose that we are given invertible 2-morphisms  $\star \lambda$  in  $\mathcal{U}(\mathfrak{sl}_2)$  for each  $\lambda \in t + 2\mathbb{Z}$ , such that  $\star \lambda$  and  $\circlearrowleft \lambda$  commute. We denote the inverse of  $\star \lambda$  by labeling the star with  $-1$ . Then there is a strict graded monoidal functor  $\Omega_t^* : \mathcal{N}(\mathcal{B}_t) \rightarrow \mathcal{U}(\mathfrak{sl}_2; t)$  taking  $B$  to  $E \oplus F$  and defined on generating morphisms by

$$\begin{aligned}
 \Omega_t^*(\bullet)_{\lambda} &:= \circlearrowleft \lambda - \circlearrowright \lambda, \\
 \Omega_t^*(\times)_{\lambda} &:= \circlearrowleft \lambda + \circlearrowright \lambda + \circlearrowleft \lambda + \circlearrowright \lambda + \circlearrowleft \lambda + \circlearrowright \lambda, \\
 \Omega_t^*(\cap)_{\lambda} &:= \circlearrowleft \lambda + \circlearrowright \lambda, \\
 \Omega_t^*(\cup)_{\lambda} &:= -\circlearrowleft \lambda + \circlearrowright \lambda
 \end{aligned}$$

for  $\lambda \in t + 2\mathbb{Z}$ . The proof of the existence of this is almost identical to the proof in Theorem 4.2—in all of the calculations stars cancel with their inverses so that the final relations that need to be checked reduce to the same ones as checked before. For example, taking  $\star \lambda := \circlearrowleft \lambda$  produces a monoidal functor  $\tilde{\Omega}_t$  with

$$\begin{aligned}
 \tilde{\Omega}_t(\uparrow)_\lambda &:= \uparrow_\lambda - \downarrow_\lambda, \\
 \tilde{\Omega}_t(\times)_\lambda &:= \begin{array}{c} \nearrow \searrow \\ \nwarrow \nearrow \end{array}_\lambda + \begin{array}{c} \nearrow \nwarrow \\ \searrow \nearrow \end{array}_\lambda + \begin{array}{c} \nearrow \nwarrow \\ \nwarrow \nearrow \end{array}_\lambda + \begin{array}{c} \nearrow \nwarrow \\ \nwarrow \nearrow \end{array}_\lambda + \begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array}_\lambda - \begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array}_\lambda - \begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array}_\lambda + \begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array}_\lambda, \\
 \tilde{\Omega}_t(\cap)_\lambda &:= \begin{array}{c} \curvearrowright \end{array}_\lambda + \begin{array}{c} \curvearrowleft \end{array}_\lambda, \\
 \tilde{\Omega}_t(\cup)_\lambda &:= \begin{array}{c} \cup \end{array}_\lambda + \begin{array}{c} \cup \end{array}_\lambda
 \end{aligned}$$

for  $\lambda \in t + 2\mathbb{Z}$ .

**Remark 4.4.** Recall from Remarks 2.2 and 3.1 that both  $\mathcal{N}_{\mathcal{B}_t}$  and  $\mathfrak{U}(\mathfrak{sl}_2)$ , hence,  $\mathfrak{u}(\mathfrak{sl}_2; t)$  can be equipped with gradings. However, the functor  $\Omega_t$  in Theorem 4.2 is *not* a graded monoidal functor—it does not preserve degrees of the generating morphisms. One way to fix this is to pass to the  $q$ -envelope  $\mathfrak{U}_q(\mathfrak{sl}_2)$  of  $\mathfrak{U}(\mathfrak{sl}_2)$  defined as in [7, Section 6] (ignoring the more complicated  $\mathbb{Z}/2$ -gradings present there). This has 1-morphisms that are formal symbols  $q^n X 1_\lambda$  for 1-morphisms  $X 1_\lambda$  in  $\mathfrak{U}(\mathfrak{sl}_2)$  and  $n \in \mathbb{Z}$ , and the 2-morphism space  $\text{Hom}_{\mathfrak{U}_q(\mathfrak{sl}_2)}(q^m X 1_\lambda, q^n Y 1_\mu)$  is  $q^{n-m} \text{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)}(X, Y)$ , where the  $q$  here is the upward<sup>2</sup> grading shift functor on the category of graded  $\mathbb{k}$ -modules  $(qV)_d := V_{d+1}$ . Horizontal and vertical composition making  $\mathfrak{U}_q(\mathfrak{sl}_2)$  into a strict graded 2-category are induced in an obvious way by the ones in  $\mathfrak{U}(\mathfrak{sl}_2)$ . Then we modify Definition 4.1, redefining  $\mathfrak{u}(\mathfrak{sl}_2; t)_{\text{ooo}}$  to be the strict graded monoidal category with objects that are words in the free monoid  $\langle \widehat{E}, F \rangle$ . For any  $X \in \langle \widehat{E}, F \rangle$  and  $\lambda \in \mathbb{Z}$ , the corresponding 1-morphism  $X 1_\lambda$  in  $\mathfrak{U}_q(\mathfrak{sl}_2)$  is defined so that  $\widehat{E} 1_\lambda := q^{-\lambda-1} E 1_\lambda$ , and then

$$\text{Hom}_{\mathfrak{u}(\mathfrak{sl}_2; t)_{\text{ooo}}}(Y, X) := \prod_{\lambda \in t+2\mathbb{Z}} \text{Hom}_{\mathfrak{U}_q(\mathfrak{sl}_2)_{\text{ooo}}}(Y 1_\lambda, X 1_\lambda)$$

for  $X, Y \in \langle \widehat{E}, F \rangle$ . The graded analog of  $\Omega_t$  can now be defined to be the graded monoidal functor given on objects by  $B \mapsto F \oplus \widehat{E}$ , and on morphisms as in Theorem 4.2; one just needs to check that this does indeed respect degrees. At the decategorified level, this modified definition of  $\Omega_t$  is consistent with the standard choice of the embedding  $\mathbf{U}_q(\mathfrak{sl}_2) \hookrightarrow \mathbf{U}_q(\mathfrak{sl}_2), B \mapsto F + qK^{-1}E$ .

### 5. THE BASIS THEOREM

Assume as before that  $\mathbb{k}$  is an integral domain in which 2 is invertible and  $t \in \{0, 1\}$ . Fix also  $m, n \geq 0$ . Any morphism  $f : B^{*n} \rightarrow B^{*m}$  is represented by a linear combination of  $m \times n$  string diagrams, i.e., string diagrams with  $m$  boundary points at the top and  $n$  boundary points at the bottom that are obtained by composing the generating morphisms from (2.1). It follows that  $\text{Hom}_{\mathcal{N}_{\mathcal{B}_t}}(B^{*n}, B^{*m})$  is 0 unless  $m \equiv n \pmod{2}$ .


The individual strings in an  $m \times n$  string diagram  $s$  are of four basic types: generalized cups (with two boundary points on the top edge), generalized caps (with two boundary points on the bottom edge), propagating strings (with one boundary point at the top and one at the bottom), and internal bubbles (no boundary points). We define an equivalence relation  $\sim$  on the set of  $m \times n$  string diagrams by declaring that  $s \sim s'$  if their strings

<sup>2</sup>If one prefers  $q$  to be the downward grading shift functor then one can instead use the  $q^{-1}$ -envelope  $\mathfrak{U}_{q^{-1}}(\mathfrak{sl}_2)$  in this place and obtain a graded monoidal functor  $\tilde{\Omega}_t$  taking  $B$  to  $F \oplus \widehat{E}$  and defined on morphisms as in Remark 4.3.

define the same matching on the set of  $m + n$  boundary points. We say that  $s$  is *reduced* if the following properties hold:

- There are no internal bubbles.
- Propagating strings have no critical points (i.e., points of slope 0).
- Generalized cups/caps (i.e., strings that connect top to top or bottom to bottom) each have exactly one critical point.
- There are no *double crossings* (i.e., two different strings which cross each other at least twice).

These assumptions imply in particular that there are no *self-intersections* (= crossings of a string with itself). For example, the first of the following undotted diagrams is not reduced (indeed, it fails all of the above conditions), while the second is reduced in the same  $\sim$ -equivalence:



Now fix a set  $\bar{D}(m, n)$  of representatives for the  $\sim$ -equivalence classes of *undotted* reduced  $m \times n$  string diagrams; the total number of such diagrams is  $(m + n - 1)!!$  if  $m \equiv n \pmod{2}$ , and there are none otherwise. For each of these  $\sim$ -equivalence class representatives, we also choose distinguished points in the interior of each of its strings that are away from points of intersection. Then let  $D(m, n)$  be the set of all morphisms  $f : B^{*n} \rightarrow B^{*m}$  obtained by closed dots labeled by some non-negative multiplicities at the elements of  $\bar{D}(m, n)$ .

Recall the commutative algebra  $\Gamma$  of Schur  $Q$ -functions and the algebra homomorphism  $\gamma_t : \Gamma \rightarrow \text{End}_{\mathcal{N}_{\mathbb{B}_t}}(\mathbb{1})$  from Corollary 2.6.

**Theorem 5.1.** *Viewing  $\text{Hom}_{\mathcal{N}_{\mathbb{B}_t}}(B^{*n}, B^{*m})$  as a  $\Gamma$ -module so that  $p \in \Gamma$  acts on  $f : B^{*n} \rightarrow B^{*m}$  by  $f \cdot p := f \star \gamma_t(p)$ , the morphism space  $\text{Hom}_{\mathcal{N}_{\mathbb{B}_t}}(B^{*n}, B^{*m})$  is free as a  $\Gamma$ -module with basis  $D(m, n)$ .*

We split the proof into two parts—spanning and linear independence.

*Proof of spanning part of Theorem 5.1.* Consider a morphism  $f \in \text{Hom}_{\mathcal{N}_{\mathbb{B}_t}}(B^{*n}, B^{*m})$  represented by an  $m \times n$  string diagram in which there is either a self-intersection or a double crossing. We claim that  $f$  can be rewritten as a linear combination of string diagrams with no self-intersections or double crossings, all of which have strictly fewer crossings than the original diagram. To see this, we can essentially ignore closed dots since, up to a sign, they can be moved freely along strings by the relations (2.5) and (2.8) modulo a linear combination of terms with strictly fewer crossings. If the diagram involves a curl  $\bigcirc$  then the morphism is 0 by the defining relations. If there are no curls, the presence of some self-intersection or double crossing implies that the diagram can be transformed using the braid relation and planar isotopy into a diagram containing a bigon  $\bigwedge$ , and the resulting morphism is 0 again by the defining relations. This last assertion is justified in the proof of [16, Theorem 8.3] by adapting an argument from [12, Lemma 2], which establishes the analogous result for closed 4-valent planar graphs (viewing a crossing in our setup as a 4-valent vertex).

Applying the claim and induction on the number of crossings, we are reduced to considering a morphism  $f$  represented by an  $m \times n$  string diagram with no self-intersections

or double crossings. Using planar isotopy, we can assume moreover that there are no cups or caps on propagating strings, and at most one cup or cap on all other strings. In particular, all of the internal bubbles are simple circles decorated by some number of dots. Using the bubble slide relation (2.27), these internal bubbles can then be moved so that they all appear on the right-hand edge of the diagram and there are no nested bubbles. Thus, we have produced a  $\Gamma$ -linear combination of morphisms defined by reduced string diagrams. In any reduced string diagram, closed dots can be moved to some distinguished point on each string modulo diagrams with fewer crossings. It remains to observe that any two undotted string diagrams of the same type are equivalent, i.e., they define the same morphism. This follows using the braid relation and planar isotopy once again. Hence, we obtain the desired  $\Gamma$ -linear combination of diagrams in  $D(m, n)$ .  $\square$

The proof of linear independence needs some additional input about the structure of the 2-category  $\mathfrak{U}(\mathfrak{sl}_2)$ . Specifically, we need bases for its 2-morphism spaces. In fact, we are going to work with a completion  $\widehat{\mathfrak{U}}(\mathfrak{sl}_2)$  of  $\mathfrak{U}(\mathfrak{sl}_2)$ . Recall from Remark 3.1 that  $\mathfrak{U}(\mathfrak{sl}_2)$  is naturally a strict graded 2-category. For 1-morphisms  $X1_\lambda, Y1_\lambda : \lambda \rightarrow \mu$  in  $\mathfrak{U}(\mathfrak{sl}_2)$  (so  $X$  and  $Y$  are words in  $\langle E, F \rangle$  of weight  $\mu - \lambda$ ), the graded 2-morphism space

$$\mathrm{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda) = \bigoplus_{d \in \mathbb{Z}} \mathrm{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda)_d \quad (5.1)$$

has finite-dimensional graded pieces and  $\mathrm{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda)_d = 0$  for  $d \ll 0$ . Consequently, it makes sense to pass to the completion with respect to the grading. This is a strict 2-category with the same objects and 1-morphisms as  $\mathfrak{U}(\mathfrak{sl}_2)$ , and 2-morphisms that are defined from

$$\mathrm{Hom}_{\widehat{\mathfrak{U}}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda) := \prod_{d \in \mathbb{Z}} \mathrm{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda)_d \quad (5.2)$$

with horizontal and vertical composition laws induced by the ones in  $\mathfrak{U}(\mathfrak{sl}_2)$ .

The *non-degeneracy* of  $\mathfrak{U}(\mathfrak{sl}_2)$  gives bases for the 2-morphism spaces  $\mathrm{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda)$  of a similar nature to the bases in Theorem 5.1. The result can be deduced from [16, 17], but it was not formulated explicitly until [15, Theorem 1.3] (which also extended the result from  $\mathfrak{sl}_2$  to  $\mathfrak{sl}_n$ ). To state the result in a way that is convenient for the present purposes, take 1-morphisms  $X1_\lambda, Y1_\lambda : \lambda \rightarrow \mu$  in  $\mathfrak{U}(\mathfrak{sl}_2)$  represented by words  $X, Y \in \langle E, F \rangle$  of lengths  $m$  and  $n$ , respectively. Take  $s \in D(n, m)$ . By writing the letters in the words  $X$  and  $Y$  at the ends of the strings at the top and bottom of the diagram  $s$ , respectively,  $s$  determines a matching between the letters of  $X$  and  $Y$ . We say that  $s$  is *admissible* for  $X$  and  $Y$  if the letters matched by each propagating string are equal and the letters matched by each generalized cup or cap are different. In that case, there is a corresponding 2-morphism  $Y1_\lambda \rightarrow X1_\lambda$  in  $\mathfrak{U}(\mathfrak{sl}_2)$  represented by the string diagram  $\vec{s} \lambda$  obtained from  $s$  by replacing the thick strings by thin strings oriented in the way dictated by the letters in  $X$  and  $Y$  ( $E$  indicates upward and  $F$  downward), replacing all closed dots with open dots with the same multiplicities, and labeling the rightmost region by  $\lambda$ . Let  $\vec{D}(Y1_\lambda, X1_\lambda)$  be the set of oriented string diagrams arising in this way from the diagrams in  $D(n, m)$  that are admissible for  $X$  and  $Y$ .

As mentioned already, the graded algebra  $\mathrm{End}_{\mathfrak{U}(\mathfrak{sl}_2)}(1_\lambda)$  can be identified with the algebra  $\Lambda$  of symmetric functions, viewed as a graded algebra so that  $e_r$  and  $h_r$  are in degree  $2r$ , so that  $\lambda \circlearrowleft(u) = u^{-\lambda}e(-u)$  and  $\lambda \circlearrowright(u) = u^\lambda h(u)$ ; this assertion is part of the non-degeneracy theorem that we are describing. Hence, each 2-morphism space

$\text{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda)$  is naturally a graded  $\Lambda$ -module,  $p \in \Lambda$  acting on  $f$  by  $f \cdot p := f \star p$ . The full theorem asserts moreover that each  $\text{Hom}_{\mathfrak{U}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda)$  is free as a graded  $\Lambda$ -module with homogeneous basis given by the set  $\vec{D}(Y1_\lambda, X1_\lambda)$ . The following is an immediate consequence.

**Theorem 5.2.** *For  $\lambda \in \mathbb{Z}$ , the endomorphism algebra  $\text{End}_{\widehat{\mathfrak{U}}(\mathfrak{sl}_2)}(1_\lambda)$  is identified with the grading completion  $\widehat{\Lambda}$  of the algebra of symmetric functions. Hence, for any 1-morphisms  $X1_\lambda, Y1_\lambda : \lambda \rightarrow \mu$ , the 2-morphism space  $\text{Hom}_{\widehat{\mathfrak{U}}(\mathfrak{sl}_2)}(Y1_\lambda, X1_\lambda)$  is naturally a topological  $\widehat{\Lambda}$ -module. In fact, it is free as a topological  $\widehat{\Lambda}$ -module with topological basis  $\vec{D}(Y1_\lambda, X1_\lambda)$ .*

We need one more basic lemma.

**Lemma 5.3.** *For any  $a \in \mathbb{k}$ , there is a strict 2-functor  $\eta_a : \mathfrak{U}(\mathfrak{sl}_2) \rightarrow \mathfrak{U}(\mathfrak{sl}_2)$  which fixes objects and 1-morphisms, and is defined on generating 2-morphisms by*

$$\begin{aligned} \uparrow \circ \lambda &\mapsto \uparrow \circ \lambda + a \uparrow \lambda, & \curvearrowright \lambda &\mapsto \curvearrowright \lambda, & \cup \lambda &\mapsto \cup \lambda, \\ \times \lambda &\mapsto \times \lambda, & \curvearrowleft \lambda &\mapsto \curvearrowleft \lambda, & \cup \lambda &\mapsto \cup \lambda. \end{aligned}$$

It also maps

$$n \circ \lambda \mapsto \sum_{r=0}^n \binom{n}{r} a^r r \circ \lambda, \quad \overset{n}{\circ} \lambda \mapsto \sum_{r=0}^n \binom{-\lambda-r}{n-r} (-a)^{n-r} \overset{r}{\circ} \lambda, \tag{5.3}$$

$$\lambda \circ n \mapsto \sum_{r=0}^n \binom{n}{r} a^r \lambda \circ r, \quad \lambda \overset{n}{\circ} \mapsto \sum_{r=0}^n \binom{\lambda-r}{n-r} (-a)^{n-r} \lambda \overset{r}{\circ} \tag{5.4}$$

for  $n \geq 0$ .

*Proof.* The existence of  $\eta_a$  follows by checking relations. All of these are clear except for (3.8), and this is easy enough to see if one works with the equivalent form (3.24). In more detail, we note that

$$\eta_a \left( \uparrow \overset{-u}{\circ} \lambda \right) = \uparrow \overset{-u-a}{\circ} \lambda,$$

because  $\eta_a(u \uparrow \lambda - \uparrow \lambda) = (u-a) \uparrow \lambda - \uparrow \lambda$ . The relation (3.24) follows using this and the observation that  $[f(u-a)]_{u-1} = [f(u)]_{u-1}$  for  $f(u) \in \mathbb{k}((u^{-1}))$ . To deduce the formulae describing  $\eta_a$  on bubbles, we explain assuming  $\lambda \geq 0$ . By the observation already made, we have that

$$\eta_a \left( \overset{u}{\circ} \lambda \right) = \overset{u-a}{\circ} \lambda.$$

Since  $\lambda \geq 0$ , we have that  $\overset{u}{\circ} \lambda = \delta_{\lambda,0} 1_{1_\lambda}$ , and it follows easily that  $\eta_a$  maps the generating function  $\lambda \circ(u)$  from (3.19) to  $\lambda \circ(u-a)$ . Inverting, we deduce that  $\eta_a$  maps  $\lambda \circ(u)$  to  $\lambda \circ(u-a)$ . The various formulae now follow by equating coefficients.  $\square$

Let  $\iota : \mathfrak{U}(\mathfrak{sl}_2) \rightarrow \widehat{\mathfrak{U}}(\mathfrak{sl}_2)$  be the canonical inclusion. The 2-functor  $\eta_a : \mathfrak{U}(\mathfrak{sl}_2) \rightarrow \mathfrak{U}(\mathfrak{sl}_2)$  maps

$$\uparrow \dots \uparrow \lambda \mapsto 2a \uparrow \uparrow \lambda + \uparrow \uparrow \lambda + \uparrow \uparrow \lambda.$$

Assuming that  $a \in \mathbb{k}^\times$ , this 2-morphism is invertible in the completion  $\widehat{\mathfrak{U}}(\mathfrak{sl}_2)$ , hence, the composition  $\iota \circ \eta_a$  extends uniquely to a strict 2-functor

$$\overline{\iota \circ \eta_a} : \mathfrak{U}(\mathfrak{sl}_2)_{\circ\dots\circ} \rightarrow \widehat{\mathfrak{U}}(\mathfrak{sl}_2). \tag{5.5}$$

Finally, for  $t \in \{0, 1\}$ , we let  $\widehat{\mathfrak{U}}(\mathfrak{sl}_2; t)$  be the strict monoidal category defined by collapsing  $\widehat{\mathfrak{U}}(\mathfrak{sl}_2)$  in exactly the same way as in Definition 4.1, replacing the localization  $\mathfrak{U}(\mathfrak{sl}_2)_{\circ\dots\circ}$  with the completion  $\widehat{\mathfrak{U}}(\mathfrak{sl}_2)$ . The 2-functor (5.5) induces a strict monoidal functor


$$\zeta_a : \mathfrak{U}(\mathfrak{sl}_2; t)_{\circ\dots\circ} \rightarrow \widehat{\mathfrak{U}}(\mathfrak{sl}_2; t). \tag{5.6}$$

We denote its natural extension to the additive envelopes of these categories by

$$\zeta_a^+ : \text{Add} \left( \mathfrak{U}(\mathfrak{sl}_2; t)_{\circ\dots\circ} \right) \rightarrow \text{Add} \left( \widehat{\mathfrak{U}}(\mathfrak{sl}_2; t) \right). \tag{5.7}$$

This functor will be useful as it is easier to work with  $\widehat{\mathfrak{U}}(\mathfrak{sl}_2; t)$  than with  $\mathfrak{U}(\mathfrak{sl}_2; t)_{\circ\dots\circ}$ , since we can exploit the topological basis arising from Theorem 5.2.

*Proof of linear independence part of Theorem 5.1.* We first use the pivotal structure to make a standard reduction: Closing diagrams on the left



defines a  $\Gamma$ -module isomorphism  $\text{Hom}_{\mathcal{N}_{\mathcal{B}_t}}(B^{*n}, B^{*m}) \xrightarrow{\sim} \text{Hom}_{\mathcal{N}_{\mathcal{B}_t}}(B^{*(m+n)}, \mathbb{1})$ . Thus, the proof is reduced to the special case that  $m = 0$ , which we assume from now on.

Take a linear relation

$$\sum_{s \in D(n,0)} s \cdot p_s = 0 \tag{5.8}$$

in  $\text{Hom}_{\mathcal{N}_{\mathcal{B}_t}}(B^{*n}, \mathbb{1})$  for  $p_s \in \Gamma$ . We must show that  $p_s = 0$  for all  $s$ . Suppose not and choose  $u \in D(n, 0)$  with a maximal number of crossings such that  $p_u \neq 0$ . Let  $Y$  be the word in  $\langle E, F \rangle$  obtained by orienting the generalized caps in  $u$  from left to right then reading the orientations of the boundary points using the usual dictionary  $E = \text{upward}$  and  $F = \text{downward}$ . Also take any  $\lambda \in \mathbb{Z}$ . We apply the monoidal functor  $\zeta_a^+ \circ \Omega_t$  to (5.8) to obtain a morphism in  $\text{Add}(\widehat{\mathfrak{U}}(\mathfrak{sl}_2; t))$ , then restrict this to  $Y1_\lambda$  to obtain a linear relation

$$\sum_{s \in D(n,0)} \zeta_a^+(\Omega_t(s \cdot p_s))|_{Y1_\lambda} = 0$$

in  $\text{Hom}_{\widehat{\mathfrak{U}}(\mathfrak{sl}_2)}(Y1_\lambda, 1_\lambda)$ . By (4.2), we deduce that

$$\sum_{s \in D(n,0)} \zeta_a^+(\Omega_t(s))|_{Y1_\lambda} \cdot \zeta_a(p_s) = 0, \tag{5.9}$$

where  $\zeta_a(p_s)$  denotes the image of  $p_s \in \Gamma \subseteq \Lambda$  under the automorphism  $\zeta_a$  of  $\Lambda \equiv \text{End}_{\mathfrak{U}(\mathfrak{sl}_2)}(1_\lambda)$  described by (5.3) and (5.4). In particular, this completes the proof in the special case  $n = 0$ .

Now we need to think more carefully about the 2-morphisms  $\zeta_a^+(\Omega_t(s))|_{Y1_\lambda}$  arising in (5.9). It is simply 0 if  $s$  is not admissible for  $Y$ . Assuming  $s$  is admissible, the definition of  $\Omega_t$  given in Theorem 4.2 implies that  $\zeta_a^+(\Omega_t(s))|_{Y1_\lambda}$  is a topological sum of morphisms defined by oriented string diagrams with the same number or with fewer crossings compared to  $s$ , with all the ones with the same number of crossings being of the same

underlying type as  $s$ ; this sum here may be infinite since the images under  $\zeta_a$  of internal bubbles and teleporters are infinite linear combinations of diagrams with extra dots and internal dotted bubbles. Using the straightening algorithm sketched in the spanning part of the proof, diagrams with the same number of crossings but a different type to  $s$  and diagrams with strictly fewer crossings than  $s$  can be rewritten as a  $\Lambda$ -linear combination of basis vectors from  $\overrightarrow{D}(Y_{1_\lambda}, 1_\lambda)$ , all of which either have the same number of crossings but a different type to  $s$  as before or have fewer crossings than  $s$ . Now let

$$X := \{s \in D(n, 0) \mid s \sim u\}.$$

For  $s \in X$ , all of the generalized caps in  $\vec{s} \lambda$  are oriented from left to right, hence, this diagram only involves upward or rightward crossings. Using the definition of  $\Omega_t$ , it follows that  $\zeta_a^+(\Omega_t(s))|_{Y_{1_\lambda}} = \zeta_a(\vec{s} \lambda) +$  (a topological linear combination of basis vectors with strictly fewer crossings). Putting these two points together, it follows that

$$\sum_{s \in D(n,0)} \zeta_a^+(\Omega_t(s))|_{Y_{1_\lambda}} \cdot \zeta_a(p_s) = \sum_{s \in X} \zeta_a(\vec{s} \lambda) \cdot \zeta_a(p_s) + (*) = 0,$$

where  $(*)$  is a topological linear combination of basis vectors with the same number of crossings as  $u$  but a different type or with strictly fewer crossings. In view of Theorem 5.2, it follows that

$$\sum_{s \in X} \zeta_a(\vec{s} \lambda) \cdot \zeta_a(p_s) = 0$$

in  $\text{Hom}_{\mathbb{U}(\mathfrak{sl}_2)}(Y_{1_\lambda}, 1_\lambda)$ , hence,  $\sum_{s \in X} (\vec{s} \lambda) \cdot p_s = 0$ . We deduce that  $p_s = 0$  for all  $s \in X$ , in particular,  $p_u = 0$ . This contradiction completes the proof of Theorem 5.1.  $\square$

**Corollary 5.4.** *The algebra homomorphism  $\gamma_t : \Gamma \rightarrow \text{End}_{\mathcal{A}(\mathbb{B}_t)}(\mathbb{1})$  is an isomorphism.*

*Proof.* This follows from the  $m = n = 0$  case of Theorem 5.1, since  $D(0, 0)$  is a singleton.  $\square$

#### ACKNOWLEDGMENTS

We thank Peng Shan for her interest and encouragement in the early stages of this project.

#### REFERENCES

- [1] Susumu Ariki, Andrew Mathas, and Hebing Rui, *Cyclotomic Nazarov–Wenzl algebras*, Nagoya Math. J. **182** (2006), 47–134.
- [2] Huanchen Bao and Weiqiang Wang, *Canonical bases arising from quantum symmetric pairs*, Invent. Math. **213** (2018), no. 3, 1099–1177.
- [3] ———, *A new approach to Kazhdan–Lusztig theory of type B via quantum symmetric pairs*, Astérisque, vol. 402, 2018.
- [4] Collin Berman and Weiqiang Wang, *Formulae of  $\nu$ -divided powers in  $U_q(\mathfrak{sl}_2)$* , J. Pure Appl. Algebra **222** (2018), no. 9, 2667–2702.
- [5] Jonathan Brundan, *On the definition of Kac–Moody 2-category*, Math. Ann. **364** (2016), 353–372.
- [6] Jonathan Brundan and Nicholas Davidson, *Categorical actions and crystals*, in Categorification and higher representation theory, Contemporary Mathematics, vol. 683, American Mathematical Society, 2017, pp. 105–147.
- [7] Jonathan Brundan and Alexander P. Ellis, *Monoidal supercategories*, Commun. Math. Phys. **351** (2017), no. 3, 1045–1089.
- [8] Jonathan Brundan, Alistair Savage, and Ben Webster, *Heisenberg and Kac–Moody categorification*, Sel. Math., New Ser. **26** (2020), Paper no. 74.
- [9] ———, *Foundations of Frobenius Heisenberg categories*, J. Algebra **578** (2021), 115–185.

- [10] ———, *The degenerate Heisenberg category and its Grothendieck ring*, *Ann. Sci. Éc. Norm. Supér.* **56** (2023), 1517–1563.
- [11] Jonathan Brundan, Ben Webster, and Weiqiang Wang, *Nil-Brauer categorifies the split  $\iota$ -quantum group of rank one*, [arXiv:2305.05877](https://arxiv.org/abs/2305.05877), 2023.
- [12] Rui Pedro Carpentier, *From planar graphs to embedded graphs—a new approach to Kauffman and Vogel’s polynomial*, *J. Knot Theory Ramifications* **9** (2000), no. 8, 975–986.
- [13] Zajj Daugherty, Arun Ram, and Rahbar Virk, *Affine and degenerate affine BMW algebras: actions on tensor space*, *Sel. Math., New Ser.* **19** (2013), no. 2, 611–653.
- [14] Michael Ehrig and Catharina Stroppel, *Nazarov–Wenzl algebras, coideal subalgebras and categorified skew Howe duality*, *Adv. Math.* **331** (2018), 58–142.
- [15] Mikhail Khovanov and Aaron D. Lauda, *A categorification of quantum  $\mathfrak{sl}(n)$* , *Quantum Topol.* **1** (2010), 1–92.
- [16] Aaron D. Lauda, *A categorification of quantum  $\mathfrak{sl}(2)$* , *Adv. Math.* **225** (2010), no. 6, 3327–3424.
- [17] ———, *Categorified quantum  $\mathfrak{sl}(2)$  and equivariant cohomology of iterated flag varieties*, *Algebr. Represent. Theory* **14** (2011), no. 2, 253–282.
- [18] Ian G. Macdonald, *Symmetric Functions and Hall Polynomials*, second ed., *Oxford Classic Texts in the Physical Sciences*, Clarendon Press, 2015.
- [19] Maxim Nazarov, *Young’s orthogonal form for Brauer’s centralizer algebra*, *J. Algebra* **182** (1996), no. 3, 664–693.
- [20] Raphaël Rouquier, *2-Kac–Moody algebras*, [arXiv:0812.5023](https://arxiv.org/abs/0812.5023), 2008.
- [21] Hebing Rui and Linliang Song, *Affine Brauer category and parabolic category  $\mathcal{O}$  in types  $B, C, D$* , *Math. Z.* **293** (2019), 503–550.

— JONATHAN BRUNDAN —

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, 1222 UNIVERSITY OF OREGON, EUGENE, OR 97403-1222, UNITED STATES

*E-mail address:* [brundan@uoregon.edu](mailto:brundan@uoregon.edu)

*URL:* <http://pages.uoregon.edu/brundan/>

— WEIQIANG WANG —

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, P. O. BOX 400137, CHARLOTTESVILLE, VA 22904-4137, UNITED STATES

*E-mail address:* [ww9c@virginia.edu](mailto:ww9c@virginia.edu)

*URL:* <https://uva.theopenscholar.com/weiqiang-wang>

— BENJAMIN WEBSTER —

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO & PERIMETER INSTITUTE FOR THEORETICAL PHYSICS, 200 UNIVERSITY AVENUE WEST, WATERLOO, ON N2L 3G1, CANADA

*E-mail address:* [ben.webster@uwaterloo.ca](mailto:ben.webster@uwaterloo.ca)

*URL:* <https://uwaterloo.ca/scholar/b2webste>